

Polynomial growth of the derivative for diffeomorphisms on tori

Krzysztof Frączek

February 1, 2008

Abstract

We consider area-preserving diffeomorphisms on tori with zero entropy. We classify ergodic area-preserving diffeomorphisms of the 3-torus for which the sequence $\{Df^n\}_{n \in \mathbb{N}}$ has polynomial growth. Roughly speaking, the main theorem says that every ergodic area-preserving C^2 -diffeomorphism with polynomial uniform growth of the derivative is C^2 -conjugate to a 2-steps skew product of the form

$$\mathbb{T}^3 \ni (x_1, x_2, x_3) \mapsto (x_1 + \alpha, \varepsilon x_2 + \beta(x_1), x_3 + \gamma(x_1, x_2)) \in \mathbb{T}^3,$$

where $\varepsilon = \pm 1$. We also indicate why there is no 4-dimensional analogue of the above result. Random diffeomorphisms on the 2-torus are studied as well.

1 Introduction

Let M be a compact Riemannian smooth manifold and let μ be a probability Borel measure with $\text{supp}(\mu) = M$. Let $f : (M, \mu) \rightarrow (M, \mu)$ be a smooth measure-preserving diffeomorphism. An important question of smooth ergodic theory is: what is the relation between asymptotic properties of the sequence $\{Df^n\}_{n \in \mathbb{N}}$ and dynamical properties of the dynamical system $f : (M, \mu) \rightarrow (M, \mu)$. There are results well describing this relation in the case where M is the torus. For example, if f is homotopic to the identity, the rotation vector of f is ergodic and the sequence $\{Df^n\}_{n \in \mathbb{N}}$ is uniformly bounded, then f is C^0 -conjugate to an ergodic rotation (see [7] p.181). Moreover, if $\{Df^n\}_{n \in \mathbb{N}}$ is bounded in the C^r -norm ($r \in \mathbb{N} \cup \{\infty\}$), then f and the ergodic rotation are C^r -conjugated (see [7] p.182). On the other hand, if $\{Df^n\}_{n \in \mathbb{N}}$ has "exponential growth", precisely if f is an Anosov diffeomorphism, then f is C^0 -conjugate to an algebraic automorphism of the torus (see [9]).

2000 *Mathematics Subject Classification*: 37A05, 37C05, 37C40.
Research partly supported by KBN grant 5 P03A 027 21(2001).

A natural question is: what can happen between the above extreme cases? The aim of this paper is to classify measure-preserving diffeomorphisms f of tori for which the sequence $\{Df^n\}_{n \in \mathbb{N}}$ has polynomial growth. One definition of the polynomial growth of the derivative is presented in [4]. In the above-mentioned paper, the following result is proved.

Proposition 1 *Let $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be an ergodic area-preserving C^2 -diffeomorphism. If the sequence $\{n^{-\tau} Df^n\}_{n \in \mathbb{N}}$ converges a.e. ($\tau > 0$) to a measurable nonzero function, then $\tau = 1$ and f is algebraically conjugate (i.e. via a group automorphism) to the skew product of an irrational rotation on the circle and a circle cocycle with nonzero topological degree.*

Moreover, in [5], the author has shown that if $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ is an ergodic area-preserving C^3 -diffeomorphism for which the sequence $\{n^{-1} Df^n\}_{n \in \mathbb{N}}$ is uniformly separated from 0 and ∞ and it is bounded in the C^2 -norm, then f is also algebraically conjugate to the skew product of an irrational rotation on the circle and a circle cocycle with nonzero topological degree.

Next very interesting result indicating how the type of growth of the derivative of a diffeomorphism reflects its important dynamical features (see [11] and [12]) says that every symplectic diffeomorphism $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ homotopic to the identity with a fixed point either is equal to the identity map or there exists $c > 0$ such that

$$\max(\|Df^n\|_{sup}, \|Df^{-n}\|_{sup}) \geq cn$$

for any natural n .

In this paper some analogous of Proposition 1 are studied. In Section 2 we discuss some random versions of Proposition 1. In Section 3 we classify area-preserving ergodic C^2 -diffeomorphisms of polynomial uniform growth of the derivative on the 3-torus, i.e. diffeomorphisms for which the sequence $\{n^{-\tau} Df^n\}_{n \in \mathbb{N}}$ converges uniformly to a non-zero function. It is shown that if the limit function is of class C^1 , then τ equals either 1 or 2 and the diffeomorphism is C^2 -conjugate to a 2-steps skew product. In Section 4 we indicate why there is no 4-dimensional analogue of the above result.

2 Random diffeomorphism on the 2-torus

Throughout this section we will consider smooth random dynamical systems over an abstract dynamical system $(\Omega, \mathcal{F}, P, T)$, where $(\Omega, \mathcal{F}, P)$ is a Lebesgue space and $T : (\Omega, \mathcal{F}, P) \rightarrow (\Omega, \mathcal{F}, P)$ is an ergodic measure-preserving automorphism. As a phase space for smooth random diffeomorphisms we will consider a compact Riemannian C^∞ -manifold M equipped with its Borel σ -algebra $\mathcal{B}$. A measurable map f

$$\mathbb{Z} \times \Omega \times M \ni (n, \omega, x) \mapsto f_\omega^n x \in M$$

satisfying for P -a.e. $\omega \in \Omega$ the following conditions

- $f_\omega^0 = \text{Id}_M$, $f_\omega^{m+n} = f_{T^n \omega}^m \circ f_\omega^n$ for all $m, n \in \mathbb{Z}$,
- $f_\omega^n : M \rightarrow M$ is a smooth function for all $n \in \mathbb{Z}$,

is called a *smooth random dynamical system* (RDS). Of course, the smooth RDS is generated by the random diffeomorphism $f_\omega = f_\omega^1$ in the sense that

$$f_\omega^n = \begin{cases} f_{T^{n-1}\omega} \circ \dots \circ f_{T\omega} \circ f_\omega & \text{for } n > 0 \\ \text{Id}_M & \text{for } n = 0 \\ f_{T^n \omega}^{-1} \circ f_{T^{n+1}\omega}^{-1} \circ \dots \circ f_{T^{-1}\omega}^{-1} & \text{for } n < 0. \end{cases}$$

Consider the skew-product transformation $T_f : (\Omega \times M, \mathcal{F} \otimes \mathcal{B}) \rightarrow (\Omega \times M, \mathcal{F} \otimes \mathcal{B})$ induced naturally by f

$$T_f(\omega, x) = (T\omega, f_\omega x).$$

Then $T_f^n(\omega, x) = (T^n \omega, f_\omega^n x)$ for all $n \in \mathbb{Z}$. We call a probability measure μ on $(\Omega \times M, \mathcal{F} \otimes \mathcal{B})$ *f-invariant*, if μ is invariant under T_f and has marginal P on Ω . Such measures can also be characterized in terms of their disintegrations μ_ω , $\omega \in \Omega$ by $f_\omega \mu_\omega = \mu_{T\omega}$ P -a.e. A f -invariant measure μ is said to be *ergodic* if $T_f : (\Omega \times M, \mathcal{F} \otimes \mathcal{B}, \mu) \rightarrow (\Omega \times M, \mathcal{F} \otimes \mathcal{B}, \mu)$ is ergodic. We say that a f -invariant measure μ has full support, if $\text{supp}(\mu_\omega) = M$ for P -a.e. $\omega \in \Omega$.

In this section we will deal with almost everywhere differentiable and C^r measure-preserving random dynamical systems with polynomial growth of the derivative. Suppose that $f : \mathbb{Z} \times \Omega \times M \rightarrow M$ is a C^0 RDS and μ is a f -invariant measure on $\Omega \times M$. The RDS f is called μ -almost everywhere differentiable if for every integer n and for μ -a.e. $(\omega, x) \in \Omega \times M$ there exists the derivative $Df_\omega^n(x) : T_x M \rightarrow T_{f_\omega^n x} M$ and

$$\int_M \|Df_\omega^n(x)\|_{n,\omega,x} d\mu_\omega(x) < \infty$$

for every $n \in \mathbb{Z}$ and P -a.e. $\omega \in \Omega$, where $\|\cdot\|_{n,\omega,x}$ is the operator norm in $\mathcal{L}(T_x M, T_{f_\omega^n x} M)$.

In the paper we will look more closely at RDS on tori. Let d be a natural number. By $\mathbb{T}^d$ we will mean the d -dimensional torus $\{(z_1, \dots, z_d) \in \mathbb{C} : |z_1| = \dots = |z_d| = 1\}$ which most often will be treated as the quotient group $\mathbb{R}^d / \mathbb{Z}^d$; $\lambda^{\otimes d}$ will denote Lebesgue measure on $\mathbb{T}^d$. We will identify functions on $\mathbb{T}^d$ with $\mathbb{Z}^d$ -periodic functions (i.e. periodic of period 1 in each coordinates) on $\mathbb{R}^d$. Let $f : \mathbb{T}^d \rightarrow \mathbb{T}^d$ be a smooth diffeomorphism. We will identify f with a diffeomorphism $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that

$$f(x_1, \dots, x_j + 1, \dots, x_d) = f(x_1, \dots, x_d) + (a_{1j}, \dots, a_{dj})$$

for every $(x_1, \dots, x_d) \in \mathbb{R}^d$, where $A = [a_{ij}]_{1 \leq i,j \leq d} \in GL_d(\mathbb{Z})$. We call A the *linear part* of the diffeomorphism f . Then there exist smooth functions $f_i : \mathbb{T}^d \rightarrow \mathbb{R}$ such that

$$f_i(x_1, \dots, x_d) = \sum_{j=1}^d a_{ij} x_j + \tilde{f}_i(x_1, \dots, x_d),$$

where $f_i : \mathbb{R}^d \rightarrow \mathbb{R}$ is the i -th coordinate functions of f .

Definition 1 We say that a μ -almost everywhere differentiable RDS f on $\mathbb{T}^d$ over $(\Omega, \mathcal{F}, P, T)$ has τ -polynomial ($\tau > 0$) growth of the derivative if

$$\frac{1}{n^\tau} Df_\omega^n(x) \rightarrow g(\omega, x) \text{ for } \mu\text{-a.e. } (\omega, x) \in \Omega \times \mathbb{T}^d,$$

where $g : \Omega \times \mathbb{T}^d \rightarrow M_d(\mathbb{R})$ is μ non-zero, i.e. there exists a set $A \in \mathcal{F} \otimes \mathcal{B}$ such that $\mu(A) > 0$ and $g(x) \neq 0$ for all $x \in A$. Moreover, if additionally Df^n belongs to $L^1((\Omega \times \mathbb{T}^d, \mu), M_d(\mathbb{R}))$ for all $n \in \mathbb{N}$ and the sequence $\{n^{-\tau} Df^n\}$ converges in $L^1((\Omega \times \mathbb{T}^d, \mu), M_d(\mathbb{R}))$ then we say that f has τ -polynomial L^1 -growth of the derivative.

We now give the example of an ergodic RDS on $\mathbb{T}^2$ with linear L^1 -growth of the derivative. For given $\tau : X \rightarrow X$, $\varphi : X \rightarrow \mathbb{T}$ and $n \in \mathbb{N}$ let

$$\varphi^{(n)} = \varphi(x) + \varphi(\tau x) + \dots + \varphi(\tau^{n-1}x).$$

Let us consider an almost everywhere differentiable RDS f on $\mathbb{T}^2$ over $(\Omega, \mathcal{F}, P, T)$ (called the random Anzai skew product) of the form

$$f_\omega(x_1, x_2) = (x_1 + \alpha(\omega), x_2 + \varphi(\omega, x_1)),$$

where the skew product $T_\alpha : (\Omega \times \mathbb{T}, P \otimes \lambda) \rightarrow (\Omega \times \mathbb{T}, P \otimes \lambda)$, $T_\alpha(\omega, x) = (T\omega, x + \alpha(\omega))$ is ergodic and $\varphi : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ is an absolutely continuous random mapping of the circle such that $D\varphi \in L^1(\Omega \times \mathbb{T}, P \otimes \lambda)$ and $\int_\Omega d(\varphi_\omega) dP\omega \neq 0$. Then the product measure $P \otimes \lambda^{\otimes 2}$ is f -invariant. The following lemma is a little bit more general version of Lemma 3 in [8].

Lemma 1 *The RDS f is ergodic and has linear L^1 -growth of the derivative.*

Proof. First note that

$$f_\omega^n(x_1, x_2) = (x_1 + \alpha^{(n)}(\omega), x_2 + \varphi^{(n)}(\omega, x_1))$$

for all $n \in \mathbb{N}$. Therefore

$$\frac{1}{n} Df_\omega^n(x_1, x_2) = \begin{bmatrix} 1/n & 0 \\ 1/n \sum_{k=0}^{n-1} D\varphi(T_\alpha^k(\omega, x_1)) & 1/n \end{bmatrix}.$$

By the ergodicity of T_α ,

$$\frac{1}{n} \sum_{k=0}^{n-1} D\varphi(T_\alpha^k(\omega, x)) \rightarrow \int_\Omega \int_{\mathbb{T}} D\varphi_\omega(y) dy dP\omega = \int_\Omega d(\varphi_\omega) dP\omega \neq 0$$

for $P \otimes \lambda$ -a.e. $(\omega, x) \in \Omega \times \mathbb{T}$ and in the L^1 -norm, which implies linear L^1 -growth of the derivatives of f .

To proof the ergodicity of f , we consider the family of unitary operators $U_m : L^2(\Omega \times \mathbb{T}, P \otimes \lambda) \rightarrow L^2(\Omega \times \mathbb{T}, P \otimes \lambda)$, $m \in \mathbb{Z}$ given by $U_m g(\omega, x) = e^{2\pi i m \varphi(\omega, x)} g(T\omega, x + \alpha(\omega))$. We now show that

$$(1) \quad \langle U_m^n g, g \rangle = \int_{\Omega \times \mathbb{T}} e^{2\pi i m \varphi^{(n)}(\omega, x)} g T_\alpha^n(\omega, x) \bar{g}(\omega, x) dP\omega dx \rightarrow 0 \text{ as } n \rightarrow \infty$$

for all $g \in L^2(\Omega \times \mathbb{T}, P \otimes \lambda)$ and $m \in \mathbb{Z} \setminus \{0\}$. Let Λ denote the set of all $g \in L^2(\Omega \times \mathbb{T}, P \otimes \lambda)$ satisfying (1). It is easy to check that Λ is a closed linear subspace of $L^2(\Omega \times \mathbb{T}, P \otimes \lambda)$. Therefore it suffices to show (1) for all functions of the form $g(\omega, x) = h(\omega) e^{2\pi i k x}$, where $h \in L^\infty(\Omega, P)$ and $k \in \mathbb{Z}$. Then

$$\begin{aligned} |\langle U_m^n g, g \rangle| &= \left| \int_{\Omega} h(T^n \omega) \bar{h}(\omega) e^{2\pi i k \alpha^{(n)}(\omega)} \left(\int_{\mathbb{T}} e^{2\pi i m \varphi^{(n)}(\omega, x)} dx \right) dP\omega \right| \\ &\leq \|h\|_{L^\infty}^2 \int_{\Omega} \left| \int_{\mathbb{T}} e^{2\pi i m \varphi^{(n)}(\omega, x)} dx \right| dP\omega. \end{aligned}$$

Let $\tilde{\varphi} : \Omega \times \mathbb{T} \rightarrow \mathbb{R}$ be an absolutely continuous random function such that $\varphi(\omega, x) = \tilde{\varphi}(\omega, x) + d(\varphi_\omega)x$. Without loss of generality we can assume that $\int_{\Omega} d(\varphi_\omega) dP\omega = a > 0$. For any natural n let

$$A_n = \{\omega \in \Omega : d(\varphi_\omega)^{(n)}/n > a/2\}.$$

By the ergodicity of T , $P(\Omega \setminus A_n) \rightarrow 0$ as $n \rightarrow \infty$. Applying integration by parts we obtain

$$\begin{aligned} \frac{1}{\|h\|_{L^\infty}^2} |\langle U_m^n g, g \rangle| &\leq P(\Omega \setminus A_n) + \int_{A_n} \left| \int_{\mathbb{T}} e^{2\pi i m \tilde{\varphi}^{(n)}(\omega, x)} d \frac{e^{2\pi i m d(\varphi_\omega)^{(n)}x}}{2\pi i m d(\varphi_\omega)^{(n)}} \right| dP\omega \\ &\leq P(\Omega \setminus A_n) + \frac{1}{\pi |m| a n} \int_{A_n} \left| \int_{\mathbb{T}} e^{2\pi i m d(\varphi_\omega)^{(n)}x} d e^{2\pi i m \tilde{\varphi}^{(n)}(\omega, x)} \right| dP\omega \\ &\leq P(\Omega \setminus A_n) + \frac{2}{\pi a n} \int_{A_n} \left| \int_{\mathbb{T}} D\tilde{\varphi}^{(n)}(\omega, x) dx \right| dP\omega \\ &\leq P(\Omega \setminus A_n) + \frac{2}{\pi a} \int_{\Omega \times \mathbb{T}} |D\tilde{\varphi}^{(n)}(\omega, x)/n| dP\omega dx. \end{aligned}$$

As $\int_{\Omega \times \mathbb{T}} D\tilde{\varphi}(\omega, x) dP\omega dx = 0$, applying Birkhoff's Ergodic Theorem for T_α we conclude that $\int_{\Omega \times \mathbb{T}} |D\tilde{\varphi}^{(n)}(\omega, x)/n| dP\omega dx$ tends to zero, which proves our claim.

Now suppose, contrary to our assertion, that f is not ergodic. Since the skew product T_α is ergodic, there exists a measurable function $g : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ and $m \in \mathbb{Z} \setminus \{0\}$ such that

$$e^{2\pi i m \varphi(\omega, x)} = g(\omega, x) \bar{g}(T_\alpha(\omega, x)).$$

Then $\langle U_m^n g, g \rangle = 1$ for all $n \in \mathbb{N}$, contrary to (1). $\square$

The aim of this section is to classify C^r random dynamical systems on the 2-torus that have polynomial (L^1) growth of the derivative and are ergodic with respect to an invariant measure of full support. We say that two random dynamical systems f and g on $\mathbb{T}^2$ over $(\Omega, \mathcal{F}, P, T)$ are algebraically conjugated if there exists a group automorphism $A : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ such that $f_\omega \circ A = A \circ g_\omega$ for P -a.e. $\omega \in \Omega$. We prove the following theorems.

Theorem 2 *Let f be a C^r random dynamical system on $\mathbb{T}^2$ over $(\Omega, \mathcal{F}, P, T)$. Let μ be an f -invariant ergodic measure of full support on $\Omega \times \mathbb{T}^2$. Suppose that f has τ -polynomial growth of the derivative. Then $\tau \geq 1$ and f is algebraically conjugate to a random skew product of the form*

$$\hat{f}_\omega(x_1, x_2) = (F_\omega(x_1), x_2 + \varphi_\omega(x_1)),$$

where $F : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ is a C^r random diffeomorphism of the circle. Moreover, if f preserves orientation, then there exist a random homeomorphism of the circle $\xi : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ and a measurable function $\alpha : \Omega \rightarrow \mathbb{T}$ such that

$$\xi_{T\omega} \circ F_\omega(x) = \xi_\omega(x) + \alpha_\omega \text{ } P\text{-a.e.}$$

and consequently f is topologically conjugate to the random skew product

$$\mathbb{T}^2 \ni (x_1, x_2) \longmapsto (x_1 + \alpha_\omega, x_2 + \varphi_\omega \circ \xi_\omega^{-1}(x_1)) \in \mathbb{T}^2.$$

Theorem 3 *Under the hypothesis of Theorem 2, if moreover f has τ -polynomial L^1 -growth of the derivative and μ is equivalent to the measure $P \otimes \lambda^{\otimes 2}$ with $d\mu/d(P \otimes \lambda^{\otimes 2}) \in L^\infty(\Omega \times \mathbb{T}^2)$, then*

- $\tau = 1$,
- *there exist a Lipschitz random diffeomorphism of the circle $\xi : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ with $D\xi, D\xi^{-1} \in L^\infty(\Omega \times \mathbb{T}, P \otimes \lambda)$ and a measurable function $\alpha : \Omega \rightarrow \mathbb{T}$ such that*

$$\xi_{T\omega} \circ F_\omega(x) = \xi_\omega(x) + \alpha_\omega \text{ } P\text{-a.e. and}$$

- $\int_\Omega d(\varphi_\omega \circ \xi_\omega^{-1}) dP\omega \neq 0$.

For convenience, the proofs of the theorems are divided into a sequence of lemmas. Let f be a C^1 random dynamical system on $\mathbb{T}^d$ over $(\Omega, \mathcal{F}, P, T)$. Let μ be a f -invariant ergodic measure of full support on $\Omega \times \mathbb{T}^d$. Suppose that f has τ -polynomial growth of the derivative. Let $g : \Omega \times \mathbb{T}^d \rightarrow M_d(\mathbb{R})$ denote the limit of the sequence $\{n^{-\tau} Df^n\}$.

Lemma 4 *For μ -a.e. $(\omega, x) \in \Omega \times \mathbb{T}^d$ and all integer n we have*

$$(2) \quad g(\omega, x) \neq 0, \quad g(\omega, x)^2 = 0 \text{ and}$$

$$(3) \quad g(\omega, x) = g(T^n \omega, f_\omega^n x) Df_\omega^n(x).$$

For $\mu \otimes \mu$ -a.e. $(\omega, x, v, y) \in \Omega \times \mathbb{T}^d \times \Omega \times \mathbb{T}^d$ we have

$$(4) \quad g(\omega, x)g(v, y) = 0 \text{ and } g(\omega, x) = Df_v(y)g(\omega, x).$$

Proof. Let $A \subset \Omega \times \mathbb{T}^d$ be a full μ -measure T_f -invariant subset such that $(\omega, x) \in A$ implies $\lim_{n \rightarrow \infty} n^{-\tau} Df_\omega^n(x) = g(\omega, x)$. Assume that $(\omega, x) \in A$. Since

$$\left(\frac{m+n}{m}\right)^\tau \frac{1}{(m+n)^\tau} Df_\omega^{m+n}(x) = \frac{1}{m^\tau} Df_{T^n \omega}^m(f_\omega^n x) Df_\omega^n(x)$$

and $(T^n \omega, f_\omega^n x) \in A$ for all $m, n \in \mathbb{N}$, letting $m \rightarrow \infty$, we obtain

$$g(\omega, x) = g(T^n \omega, f_\omega^n x) Df_\omega^n(x) \quad \text{for } (\omega, x) \in A \text{ and } n \in \mathbb{N}.$$

Let $B = \{(\omega, x) \in A : g(\omega, x) \neq 0\}$. By the above, B is T_f -invariant. As g is μ non-zero, we have $\mu(B) = 1$, by the ergodicity of T_f .

By the Jewett-Krieger Theorem, we can assume that Ω is a compact metric space, $T : \Omega \rightarrow \Omega$ is an uniquely ergodic homeomorphism and P is the unique T -invariant measure. Now choose a sequence $\{A_k\}_{k \in \mathbb{N}}$ of measurable subsets of A such that the functions $g, Df : A_k \rightarrow M_d(\mathbb{R})$ are continuous, all non-empty open subsets of A_k (in the induced topology) have positive measure and $\mu(A_k) > 1 - 1/k$ for any natural k . Since the transformation $(T_f)_{A_k} : (A_k, \mu_{A_k}) \rightarrow (A_k, \mu_{A_k})$ induced by T_f on A_k is ergodic, for every natural k we can find a measurable subset $B_k \subset A_k$ such that for any $(\omega, x) \in B_k$ the orbit $\{(T_f)_{A_k}^n(\omega, x)\}_{n \in \mathbb{N}}$ is dense in A_k in the induced topology and $\mu(B_k) = \mu(A_k)$.

Let $(\omega, x), (v, y) \in B_k$. Then there exists an increasing sequence $\{m_i\}_{i \in \mathbb{N}}$ of natural numbers such that $(T_f)_{A_k}^{m_i}(\omega, x) \rightarrow (v, y)$. Hence there exists an increasing sequence $\{n_i\}_{i \in \mathbb{N}}$ of natural numbers such that $T_f^{n_i}(\omega, x) \rightarrow (v, y)$ and $T_f^{n_i}(\omega, x) \in A_k$ for all $i \in \mathbb{N}$. Since $g, Df : A_k \rightarrow M_d(\mathbb{R})$ are continuous, we get $g(T^{n_i} \omega, f_\omega^{n_i} x) \rightarrow g(v, y)$ and $Df_{T^{n_i} \omega}(f_\omega^{n_i} x) \rightarrow Df_v(y)$. Since

$$\frac{1}{n_i^\tau} g(\omega, x) = g(T^{n_i} \omega, f_\omega^{n_i} x) \frac{1}{n_i^\tau} Df_\omega^{n_i}(x),$$

letting $i \rightarrow \infty$, we obtain $g(v, y) g(\omega, x) = 0$. Since

$$\frac{1}{n_i^\tau} Df_\omega^{n_i+1}(x) = Df_{T^{n_i} \omega}(f_\omega^{n_i} x) \frac{1}{n_i^\tau} Df_\omega^{n_i}(x),$$

letting $i \rightarrow \infty$, we obtain $g(\omega, x) = Df_v(y) g(\omega, x)$. Therefore

$$\mu \otimes \mu \{(\omega, x, v, y) \in \Omega \times \mathbb{T}^d \times \Omega \times \mathbb{T}^d : g(v, y) g(\omega, x) = 0\} > (1 - \frac{1}{k})^2,$$

$$\mu \{(\omega, x) \in \Omega \times \mathbb{T}^d : g(\omega, x)^2 = 0\} > 1 - \frac{1}{k}$$

and

$$\mu \otimes \mu \{(\omega, x, v, y) \in \Omega \times \mathbb{T}^d \times \Omega \times \mathbb{T}^d : g(\omega, x) = Df_v(y) g(\omega, x)\} > (1 - \frac{1}{k})^2$$

for any natural k , which proves the lemma. $\square$

Let us return to the case where $d = 2$. Suppose that A, B are non-zero real 2×2 -matrixes such that $A^2 = B^2 = AB = 0$. Then (see Lemma 4 in [4]) there exist real numbers $a, b \neq 0$ and c such that

$$A = a \begin{bmatrix} c \\ 1 \end{bmatrix} \begin{bmatrix} 1 & -c \end{bmatrix} \quad \text{and} \quad B = b \begin{bmatrix} c \\ 1 \end{bmatrix} \begin{bmatrix} 1 & -c \end{bmatrix}$$

or

$$A = a \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} \quad \text{and} \quad B = b \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix}$$

It follows that g can be represented as follows

$$g = h \begin{bmatrix} c \\ 1 \end{bmatrix} \begin{bmatrix} 1 & -c \end{bmatrix},$$

where $h : \Omega \times \mathbb{T}^2 \rightarrow \mathbb{R}$ is a measurable function, which is non-zero at μ -a.e. point and $c \in \mathbb{R}$. We can omit the second case where

$$g = h \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix},$$

because it reduces to case $c = 0$ after interchanging the coordinates, which is an algebraic isomorphism. Then from (4) we obtain

$$(5) \quad \begin{bmatrix} c \\ 1 \end{bmatrix} = Df_\omega x \begin{bmatrix} c \\ 1 \end{bmatrix}$$

for P -a.e. $\omega \in \Omega$ and for all $x \in \mathbb{T}^2$, because μ has full support. From (3) we obtain

$$(6) \quad h(\omega, x) \begin{bmatrix} 1 & -c \end{bmatrix} = h(T\omega, f_\omega x) \begin{bmatrix} 1 & -c \end{bmatrix} Df_\omega x$$

for μ -a.e. $(\omega, x) \in \Omega \times \mathbb{T}^2$.

Lemma 5 *If c is irrational, then $f_\omega(x_1, x_2) = (x_1 + \alpha(\omega), x_2 + \gamma(\omega))$, where $\alpha, \gamma : \Omega \rightarrow \mathbb{T}$ are measurable functions. Consequently, the sequence $n^{-\tau} Df^n$ tends uniformly to zero.*

Proof. From (5) we have

$$c = c \frac{\partial(f_\omega)_1}{\partial x_1} + \frac{\partial(f_\omega)_1}{\partial x_2} \quad \text{and} \quad 1 = c \frac{\partial(f_\omega)_2}{\partial x_1} + \frac{\partial(f_\omega)_2}{\partial x_2}$$

for P -a.e. $\omega \in \Omega$. It follows that for $i = 1, 2$ there exists a C^{r+1} random function $u_i : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f_i(\omega, x_1, x_2) = x_i + u_i(\omega, x_1 - cx_2).$$

Represent f as follows

$$\begin{aligned} f_1(\omega, x_1, x_2) &= a_{11}(\omega)x_1 + a_{12}(\omega)x_2 + \tilde{f}_1(\omega, x_1, x_2), \\ f_2(\omega, x_1, x_2) &= a_{21}(\omega)x_1 + a_{22}(\omega)x_2 + \tilde{f}_2(\omega, x_1, x_2), \end{aligned}$$

where $\{a_{ij}(\omega)\}_{i,j=1,2} \in GL_2(\mathbb{Z})$ and $\tilde{f}_1, \tilde{f}_2 : \Omega \times \mathbb{T}^2 \rightarrow \mathbb{R}$. Then

$$u_1(\omega, x+1) = (a_{11}(\omega) - 1)(x+1) + \tilde{f}_1(\omega, x+1, 0) = u_1(\omega, x) + a_{11}(\omega) - 1$$

and

$$u_1(\omega, x+c) = (a_{11}(\omega) - 1)x - a_{12}(\omega) + \tilde{f}_1(\omega, x, -1) = u_1(\omega, x) - a_{12}(\omega).$$

Therefore

$$a_{11}(\omega) - 1 = \lim_{x \rightarrow +\infty} \frac{u_1(\omega, x)}{x} = -a_{12}(\omega)/c$$

for μ -a.e. $\omega \in \Omega$. Since c is irrational, we conclude that $a_{11}(\omega) - 1 = a_{12}(\omega) = 0$, hence that $u_1(\omega, \cdot)$ is 1 and c periodic, and finally that $u_1(\omega, \cdot)$ is a constant function for μ -a.e. $\omega \in \Omega$. It is clear that the same conclusion can be drawn for u_2 , which completes the proof. $\square$

Lemma 6 *If c is rational, then there exist a group automorphism $A : \mathbb{T}^2 \rightarrow \mathbb{T}^2$, a C^r random diffeomorphism of the circle $F : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ and a C^r random function $\varphi : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ such that*

$$A \circ f_\omega \circ A^{-1}(x_1, x_2) = (F_\omega x_1, x_2 + \varphi_\omega(x_1)).$$

Moreover,

$$(7) \quad h_{T\omega} \circ A^{-1}(F_\omega(x_1), x_2 + \varphi_\omega(x_1)) D F_\omega(x_1) = h_\omega \circ A^{-1}(x_1, x_2)$$

for $\hat{\mu}$ -a.e. $(\omega, x_1, x_2) \in \Omega \times \mathbb{T}^2$, where $\hat{\mu} := (\text{Id}_\Omega \times A)\mu$ and $h_\omega \circ A^{-1} : \mathbb{T}^2 \rightarrow \mathbb{R}$ depends only on the first coordinate.

Proof. Let p and q be integer numbers such that $q > 0$, $\gcd(p, q) = 1$ and $c = p/q$. Choose $a, b \in \mathbb{Z}$ with $ap - bq = 1$. Consider the group automorphism $A : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ associated to the matrix $A = \begin{bmatrix} q & -p \\ -b & a \end{bmatrix}$. Then $A^{-1} = \begin{bmatrix} a & p \\ b & q \end{bmatrix}$. Let us consider the C^r RDS $\hat{f}_\omega = A \circ f_\omega \circ A^{-1}$. Then $\hat{\mu}$ is a $\hat{f}$ -invariant measure and

$$D\hat{f}_\omega(x) = A Df_\omega(A^{-1}x) A^{-1}.$$

From (5) we see that

$$\begin{bmatrix} p \\ q \end{bmatrix} = Df_\omega x \begin{bmatrix} p \\ q \end{bmatrix}$$

for P -a.e. $\omega \in \Omega$ and all $x \in \mathbb{T}^2$, hence that

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} = D\hat{f}_\omega x \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

for P -a.e. $\omega \in \Omega$ and all $x \in \mathbb{T}^2$. From (6) we see that

$$h_\omega(x) \begin{bmatrix} q & -p \end{bmatrix} = h_{T\omega}(f_\omega x) \begin{bmatrix} q & -p \end{bmatrix} Df_\omega x$$

for μ -a.e. $(\omega, x) \in \Omega \times \mathbb{T}^2$, hence that

$$h_\omega \circ A^{-1}(x) \begin{bmatrix} 1 & 0 \end{bmatrix} = h_{T\omega} \circ A^{-1}(\hat{f}_\omega x) \begin{bmatrix} 1 & 0 \end{bmatrix} D\hat{f}_\omega x$$

for $\hat{\mu}$ -a.e. $(\omega, x) \in \Omega \times \mathbb{T}^2$. It follows that

$$\frac{\partial(\hat{f}_\omega)_1}{\partial x_2} = 0, \quad \frac{\partial(\hat{f}_\omega)_2}{\partial x_2} = 1$$

for P -a.e. $\omega \in \Omega$ and

$$h_{T\omega} \circ A^{-1} \circ \hat{f}_\omega \frac{\partial(\hat{f}_\omega)_1}{\partial x_1} = h_{T\omega} \circ A^{-1}$$

for $\hat{\mu}$ -a.e. $(\omega, x) \in \Omega \times \mathbb{T}^2$. Therefore

$$\hat{f}_\omega(x_1, x_2) = (F_\omega x_1, x_2 + \varphi_\omega(x_1)),$$

where $F, \varphi : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ are C^r random functions and

$$h_{T\omega} \circ A^{-1}(F_\omega(x_1), x_2 + \varphi_\omega(x_1)) DF_\omega(x_1) = h_\omega \circ A^{-1}(x_1, x_2)$$

for $\hat{\mu}$ -a.e. $(\omega, x_1, x_2) \in \Omega \times \mathbb{T}^2$. Since $\hat{f}_\omega : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ is a C^r -diffeomorphism, we conclude that $F_\omega : \mathbb{T} \rightarrow \mathbb{T}$ is a C^r -diffeomorphism for P -a.e. $\omega \in \Omega$. Since

$$\frac{1}{n^\tau} Df_\omega^n(x) \rightarrow h_\omega(x) \begin{bmatrix} p/q \\ 1 \end{bmatrix} \begin{bmatrix} 1 & -p/q \end{bmatrix}$$

for μ -a.e. $(\omega, x_1, x_2) \in \Omega \times \mathbb{T}^2$, we have

$$\frac{1}{n^\tau} D\hat{f}_\omega^n(x) \rightarrow h_\omega(A^{-1}x)/q^2 \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

for $\hat{\mu}$ -a.e. $(\omega, x_1, x_2) \in \Omega \times \mathbb{T}^2$. Let $\hat{h}_\omega := h_\omega \circ A^{-1}$. Then

$$\frac{1}{n^\tau} \sum_{k=0}^{n-1} D\varphi_{T^k\omega}(F_\omega^k(x_1)) DF_\omega^k(x_1) \rightarrow \hat{h}_\omega(x_1, x_2)/q^2$$

for $\hat{\mu}$ -a.e. $(\omega, x_1, x_2) \in \Omega \times \mathbb{T}^2$. It follows that $\hat{h}_\omega$ depends only on the first coordinate. $\square$

Proof of Theorem 2. By Lemmas 5 and 6, to prove the first claim of the theorem we only need to show that $\tau \geq 1$. Suppose that $\tau < 1$. Let $\nu := (\text{Id}_\Omega \times \pi)\hat{\mu}$, where $\pi : \mathbb{T}^2 \rightarrow \mathbb{T}$ is the projection on the first coordinate. Then ν is a F -invariant ergodic measure of full support on $\Omega \times \mathbb{T}$. By Lemma 6,

$$\hat{h}_{T^k\omega}(F_\omega^k(x)) DF_\omega^k(x) = \hat{h}_\omega(x)$$

and

$$(8) \quad \frac{1}{n^\tau} \sum_{k=0}^{n-1} D\varphi_{T^k\omega}(F_\omega^k(x)) DF_\omega^k(x) \rightarrow \hat{h}_\omega(x)/q^2$$

for ν -a.e. $(\omega, x) \in \Omega \times \mathbb{T}$. Therefore

$$(9) \quad \frac{1}{n^\tau} \sum_{k=0}^{n-1} D\varphi_{T^k\omega}(F_\omega^k(x))/\hat{h}_{T^k\omega}(F_\omega^k(x)) \rightarrow 1/q^2$$

and consequently

$$\frac{1}{n} \sum_{k=0}^{n-1} D\varphi_{T^k\omega}(F_\omega^k(x))/\hat{h}_{T^k\omega}(F_\omega^k(x)) \rightarrow 0$$

for ν -a.e. $(\omega, x) \in \Omega \times \mathbb{T}$. It follows that the measurable cocycle $D\varphi/\hat{h} : \Omega \times \mathbb{T} \rightarrow \mathbb{R}$ over the skew product T_F is recurrent (see [13]). Consequently, for ν -a.e. $(\omega, x) \in \Omega \times \mathbb{T}$ there exists an increasing sequence of natural numbers $\{n_i\}_{i \in \mathbb{N}}$ such that

$$\left| \sum_{k=0}^{n_i-1} D\varphi_{T^k\omega}(F_\omega^k(x))/\hat{h}_{T^k\omega}(F_\omega^k(x)) \right| \leq 1.$$

It follows that

$$\frac{1}{n_i^\tau} \sum_{k=0}^{n_i-1} D\varphi_{T^k\omega}(F_\omega^k(x))/\hat{h}_{T^k\omega}(F_\omega^k(x)) \rightarrow 0,$$

contrary to (9).

Now suppose that f preserves orientation. Let us decompose $\nu_\omega = \nu_\omega^d + \nu_\omega^c$, where ν_ω^d is the discrete and ν_ω^c is the continuous part of the measure ν_ω . As this decomposition is measurable we can consider the measures $\nu^d = \int_\Omega \nu_\omega^d dP\omega$ and $\nu^c = \int_\Omega \nu_\omega^c dP\omega$ on $\Omega \times \mathbb{T}$. It is easy to check that ν^d and ν^c are F -invariant. By the ergodicity of ν , either $\nu = \nu^d$ or $\nu = \nu^c$.

We now show that $\nu = \nu^c$. Suppose, contrary to our claim, that $\nu = \nu^d$. Let $\Delta : \Omega \times \mathbb{T} \rightarrow [0, 1]$ denote the measurable function given by $\Delta(\omega, x) = \nu_\omega(\{x\})$. As ν is F -invariant we have

$$\Delta(T\omega, F_\omega x) = \nu_{T\omega}(\{F_\omega x\}) = F_\omega^{-1} \nu_{T\omega}(\{x\}) = \nu_\omega(\{x\}) = \Delta(\omega, x)$$

and consequently Δ is T_F -invariant. By the ergodicity of T_F , the function Δ is ν constant. It follows that the measure ν_ω has only finite number of atoms for P -a.e. $\omega \in \Omega$, which contradicts the fact that ν has full support.

Define $\xi_\omega(x) := \int_0^x d\nu_\omega$ for all $x \in \mathbb{R}$. Then $\xi_\omega(x+1) = \xi_\omega(x) + 1$, because $\int_x^{x+1} d\nu_\omega = 1$. Since ν_ω is continuous and ν has full support, the function $\xi_\omega : \mathbb{R} \rightarrow \mathbb{R}$ is continuous and strictly increasing. Therefore $\xi : \Omega \times \mathbb{T} \rightarrow \mathbb{T}$ is a random homeomorphism. As ν is F -invariant and F preserves orientation we have

$$\xi_{T\omega}(F_\omega x) = \int_0^{F_\omega x} d\nu_{T\omega} = \int_0^{F_\omega 0} d\nu_{T\omega} + \int_{F_\omega 0}^{F_\omega x} dF_\omega \nu_\omega = \alpha_\omega + \int_0^x d\nu_\omega = \xi_\omega(x) + \alpha_\omega$$

for P -a.e. $\omega \in \Omega$, where $\alpha_\omega = \int_0^{F_\omega 0} d\nu_{T_\omega}$. $\square$

Proof of Theorem 3 Suppose that f has τ -polynomial L^1 -growth of the derivative and μ is equivalent to $P \otimes \lambda^{\otimes 2}$. Then $DF, D\varphi \in L^1(\Omega \times \mathbb{T}, \nu)$ and $\hat{\mu}$ is equivalent to $P \otimes \lambda^{\otimes 2}$. Let $\theta \in L^1(\Omega \times \mathbb{T}^2, P \otimes \lambda^{\otimes 2})$ denote the Radon–Nikodym derivative of $\hat{\mu}$ with respect to $P \otimes \lambda^{\otimes 2}$. Then

$$\theta_{T_\omega}(F_\omega(x_1), x_2 + \varphi_\omega(x_1)) DF_\omega(x_1) = \theta_\omega(x_1, x_2)$$

for $P \otimes \lambda^{\otimes 2}$ -a.e. $(\omega, x_1, x_2) \in \Omega \times \mathbb{T}^2$. From (7) there exists a non-zero constant C such that $\theta_\omega(x_1, x_2) = Ch_\omega(x_1)$ for $P \otimes \lambda^{\otimes 2}$ -a.e. $(\omega, x_1, x_2) \in \Omega \times \mathbb{T}^2$. Then the random homeomorphism $\xi_\omega : \mathbb{T} \rightarrow \mathbb{T}$ given by $\xi_\omega(x) := \int_0^x d\nu_\omega = \int_0^x \theta_\omega(t) dt$ is a Lipschitz random diffeomorphism, because θ and $1/\theta$ are bounded. It follows that f is Lipschitz conjugate to the random skew product

$$(T_{\alpha, \psi})_\omega(x_1, x_2) = (x_1 + \alpha_\omega, x_2 + \psi_\omega(x_1)),$$

where $\psi_\omega := \varphi_\omega \circ \xi_\omega^{-1}$. From (8) we conclude that $T_{\alpha, \psi}$ has τ -polynomial L^1 -growth of the derivative and

$$\frac{1}{n^\tau} \sum_{k=0}^{n-1} D\psi(T_\alpha(\omega, x)) \rightarrow \hat{h}_\omega \circ \xi_\omega^{-1}(x) D\xi_\omega^{-1}(x)/q^2$$

in $L^1(\Omega \times \mathbb{T}, P \otimes \lambda)$. It follows that

$$n^{1-\tau} \int_\Omega d(\psi_\omega) dP_\omega \rightarrow 1/Cq^2,$$

and finally that $\tau = 1$ and $\int_\Omega d(\psi_\omega) dP_\omega \neq 0$. $\square$

3 Area-preserving diffeomorphisms of the 3-torus

In this section we give a classification of area-preserving ergodic diffeomorphisms of polynomial uniform growth of the derivative on the 3-torus, i.e. diffeomorphisms of polynomial growth of the derivative for which the sequence $\{n^{-\tau} Df^n\}_{n \in \mathbb{N}}$ converges uniformly. We first give a sequence of essential examples of such diffeomorphisms. We will consider 2-steps skew products $T_{\alpha, \beta, \gamma} : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ given by

$$T_{\alpha, \beta, \gamma}(x_1, x_2, x_3) = (x_1 + \alpha, x_2 + \beta(x_1), x_3 + \gamma(x_1, x_2)),$$

where α is irrational and $\beta : \mathbb{T} \rightarrow \mathbb{T}$, $\gamma : \mathbb{T}^2 \rightarrow \mathbb{T}$ are of class C^1 . We will denote by $d_i(\gamma)$ the topological degree of γ with respect to the i -th coordinate for $i = 1, 2$. We will use the symbol h_{x_1} to denote the partial derivative $\partial h / \partial x_1$.

Example 1. Assume that β is a constant function, $\alpha, \beta, 1$ are rationally independent and $(d_1(\gamma), d_2(\gamma)) \neq 0$. Then

$$\frac{1}{n}DT_{\alpha, \beta, \gamma}^n \rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ d_1(\gamma) & d_2(\gamma) & 0 \end{bmatrix} \neq 0$$

uniformly and $T_{\alpha, \beta, \gamma}$ is ergodic, by Lemma 1.

Example 2. Assume that γ depends only on the first coordinate, $d(\beta) = 0$, $d(\gamma) \neq 0$ and the skew product

$$\mathbb{T}^2 \ni (x_1, x_2) \mapsto (x_1 + \alpha, x_2 + \beta(x_1)) \in \mathbb{T}^2$$

is ergodic. Then

$$\frac{1}{n}DT_{\alpha, \beta, \gamma}^n \rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ d_1(\gamma) & 0 & 0 \end{bmatrix} \neq 0$$

uniformly. To prove the ergodicity $T_{\alpha, \beta, \gamma}$, it suffices to show the ergodicity of the skew product

$$\mathbb{T}^2 \ni (x_1, x_2) \mapsto (x_1 + \alpha, x_2 + m_1\beta(x_1) + m_2\gamma(x_1)) \in \mathbb{T}^2$$

for every $(m_1, m_2) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$. If $m_2 = 0$, then the ergodicity of the skew product follows from assumption. Otherwise, the claim follows from Lemma 1.

Example 3. Assume that $d(\beta) \neq 0$ and $d_2(\gamma) \neq 0$. By Lemma 1, $T_{\alpha, \beta, \gamma}$ is ergodic. We now show that $T_{\alpha, \beta, \gamma}$ has square uniform growth of the derivative. Precisely, we show that

$$(10) \quad \frac{1}{n^2}DT_{\alpha, \beta, \gamma}^n \rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ d(\beta)d_2(\gamma)/2 & 0 & 0 \end{bmatrix} \neq 0$$

uniformly. Let

$$T_\alpha(x) = x + \alpha \text{ and } T_{\alpha, \beta}(x_1, x_2) = (x_1 + \alpha, x_2 + \beta(x_1)).$$

Note that

$$T_{\alpha, \beta, \gamma}^n(x_1, x_2, x_3) = (x_1 + n\alpha, x_2 + \beta^{(n)}(x_1), x_3 + \gamma^{(n)}(x_1, x_2)).$$

Therefore

$$\begin{aligned} & \frac{1}{n^2}DT_{\alpha, \beta, \gamma}^n(x_1, x_2, x_3) \\ &= \begin{bmatrix} n^{-2} & 0 & 0 \\ n^{-2}(D\beta)^{(n)}(x_1) & n^{-2} & 0 \\ n^{-2}(\gamma_{x_1}^{(n)}(x_1, x_2) + \sum_{k=0}^{n-1} \gamma_{x_2}(T_{\alpha, \beta}^k(x_1, x_2))(D\beta)^{(k)}(x_1)) & n^{-2}(\gamma_{x_2})^{(n)}(x_1, x_2) & n^{-2} \end{bmatrix} \end{aligned}$$

To obtain (10), it only remains to verify

$$\frac{1}{n^2} \sum_{k=0}^{n-1} \gamma_{x_2}(T_{\alpha,\beta}^k(x_1, x_2))(D\beta)^{(k)}(x_1) \rightarrow d(\beta)d_2(\gamma)/2 \neq 0$$

uniformly. First observe that

$$\frac{1}{n^2} \sum_{k=0}^{n-1} (n-k) \gamma_{x_2}(T_{\alpha,\beta}^k(x_1, x_2)) = \frac{1}{n^2} \sum_{k=1}^n (\gamma_{x_2})^{(k)}(x_1, x_2) \rightarrow d_2(\gamma)/2$$

uniformly. Indeed, set $a_n := \sum_{k=1}^n (\gamma_{x_2})^{(k)}$ and $b_n := n^2$. Then

$$\frac{a_{n+1} - a_n}{b_{n+1} - b_n} = \frac{\sum_{k=0}^{n-1} \gamma_{x_2} \circ T_{\alpha,\beta}^k}{2n+1} \rightarrow \frac{1}{2} \int_{\mathbb{T}^2} \gamma_{x_2}(x_1, x_2) dx_1 dx_2 = d_2(\gamma)/2$$

uniformly. It follows that $a_n/b_n \rightarrow d(\beta)d_2(\gamma)/2$ uniformly. Hence

$$\frac{1}{n^2} \sum_{k=0}^{n-1} k \gamma_{x_2} \circ T_{\alpha,\beta}^k = \frac{1}{n} \sum_{k=0}^{n-1} \gamma_{x_2} \circ T_{\alpha,\beta}^k - \frac{1}{n^2} \sum_{k=0}^{n-1} (n-k) \gamma_{x_2} \circ T_{\alpha,\beta}^k \rightarrow d_2(\gamma)/2$$

uniformly. Then

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left\| \frac{1}{n^2} \sum_{k=0}^{n-1} \gamma_{x_2} \circ T_{\alpha,\beta}^k \cdot D\beta^{(k)} - d(\beta)d_2(\gamma)/2 \right\|_{\sup} \\ &= \lim_{n \rightarrow \infty} \left\| \frac{1}{n^2} \sum_{k=0}^{n-1} \gamma_{x_2} \circ T_{\alpha,\beta}^k \cdot D\beta^{(k)} - \frac{1}{n^2} \sum_{k=0}^{n-1} k d(\beta) \gamma_{x_2} \circ T_{\alpha,\beta}^k \right\|_{\sup} \\ &\leq \lim_{n \rightarrow \infty} \frac{\|\gamma\|_{C^1}}{n} \sum_{k=0}^{n-1} \left\| \frac{1}{k} \sum_{j=0}^{k-1} D\beta \circ T_{\alpha}^j - d(\beta) \right\|_{\sup} \\ &\leq \lim_{n \rightarrow \infty} \frac{\|\gamma\|_{C^1}}{n} \sum_{k=0}^{n-1} \left\| \frac{1}{k} \sum_{j=0}^{k-1} D\beta \circ T_{\alpha}^j - \int_{\mathbb{T}} D\beta(x) dx \right\|_{\sup} = 0. \end{aligned}$$

Example 4. Assume that

$$f(x_1, x_2, x_3) = (x_1 + \alpha, -x_2 + \beta(x_1), x_3 + \gamma(x_1)),$$

where α is irrational and $\beta, \gamma : \mathbb{T} \rightarrow \mathbb{T}$ are of class C^1 . Suppose that $d(\gamma) \neq 0$ and the factor map

$$\hat{f} : \mathbb{T}^2 \rightarrow \mathbb{T}^2, \quad \hat{f}(x_1, x_2) = (x_1 + \alpha, -x_2 + \beta(x_1))$$

is ergodic. It is easy to see that $\hat{f}$ is ergodic iff $\hat{f}^2(x_1, x_2) = (x_1 + 2\alpha, x_2 - \beta(x_1) + \beta(x_1 + \alpha))$ is. Since $d(-\beta + \beta \circ T_{\alpha}) = 0$ and $d(\gamma + \gamma \circ T_{\alpha}) \neq 0$, we conclude the diffeomorphism

$$f^2(x_1, x_2, x_3) = (x_1 + 2\alpha, x_2 - \beta(x_1) + \beta(x_1 + \alpha), x_3 + \gamma(x_1) + \gamma(x_1 + \alpha))$$

is ergodic, by Example 2. Consequently, f is ergodic. Moreover

$$\frac{1}{n}Df^n \rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ d(\gamma) & 0 & 0 \end{bmatrix} \neq 0$$

uniformly.

Example 5. Assume that an area-preserving C^2 -diffeomorphism $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ has τ -polynomial uniform growth of the derivative and

$$\frac{1}{n^\tau}Df^n \rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ h_1 & h_2 & 0 \end{bmatrix} \neq 0,$$

where $h = (h_1, h_2) : \mathbb{T}^3 \rightarrow \mathbb{R}^2$ is of class C^1 . Let $\varphi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be an area-preserving C^2 -diffeomorphism. Let $\hat{f} : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ denote the area-preserving diffeomorphism that is conjugate to f via the diffeomorphism $\varphi \times \text{Id} : \mathbb{T}^3 \rightarrow \mathbb{T}^3$, i.e.

$$\hat{f} = (\varphi \times \text{Id})^{-1} \circ f \circ (\varphi \times \text{Id}).$$

Then

$$\frac{1}{n^\tau}D\hat{f}^n(x_1, x_2, x_3) \rightarrow \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ Dh(\varphi(x_1, x_2), x_3)D\varphi(x_1, x_2) & 0 \end{bmatrix} \neq 0$$

uniformly. It follows that $\hat{f}$ has τ -polynomial uniform growth of the derivative and the limit function $\lim_{n \rightarrow \infty} n^{-\tau}D\hat{f}^n$ is of class C^1 .

In this section we prove the following theorem.

Theorem 7 *Let $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ be an area-preserving ergodic C^2 -diffeomorphism with τ -polynomial uniform growth of the derivative ($\tau > 0$). Suppose that the limit function $\lim_{n \rightarrow \infty} n^{-\tau}Df^n$ is of class C^1 . Then either $\tau = 1$ or $\tau = 2$ and f is C^2 -conjugate to a diffeomorphism of the form*

$$\mathbb{T}^3 \ni (x_1, x_2, x_3) \mapsto (x_1 + \alpha, \varepsilon x_2 + \beta(x_1), x_3 + \gamma(x_1, x_2)) \in \mathbb{T}^3,$$

where $\varepsilon = \det Df = \pm 1$.

As in the previous section, the proof of the main theorem is divided into a sequence of lemmas. Suppose that $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ is an area-preserving ergodic diffeomorphism with τ -polynomial growth of the derivative. Let $g : \mathbb{T}^3 \rightarrow M_3(\mathbb{R})$ denote the limit of the sequence $\{n^{-\tau}Df^n\}_{n \in \mathbb{N}}$. By Lemma 4, $g(\bar{x})g(\bar{y}) = 0$ and $g(\bar{x})^2 = 0$ for all $\bar{x}, \bar{y} \in \mathbb{T}^3$.

Lemma 8 *Suppose that A, B are non-zero real 3×3 -matrixes such that $A^2 = B^2 = AB = BA = 0$. Then there exist three non-zero vectors (real 1×3 -matrixes) $\bar{a}, \bar{b}, \bar{c}$ such that*

- $A = \bar{a}^T \bar{b}$ and $B = \bar{a}^T \bar{c}$ where $\bar{b}\bar{a}^T = 0$ and $\bar{c}\bar{a}^T = 0$ or
- $A = \bar{a}^T \bar{c}$ and $B = \bar{b}^T \bar{c}$ where $\bar{c}\bar{a}^T = 0$ and $\bar{c}\bar{b}^T = 0$.

Proof. Suppose that $\lambda \in \mathbb{C}$ is an eigenvalue of A for an eigenvector $\bar{x} \in \mathbb{C}^3$. Then $\lambda^2 \bar{x} = A^2 \bar{x} = 0$ and consequently $\lambda = 0$. It follows that Jordan canonical form of A equals either

$$\begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \text{ or } \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

But the second case can not occur since

$$\begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}^2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \neq 0.$$

It follows that there exists $C \in GL_3(\mathbb{R})$ such that

$$A = C \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} C^{-1} = \begin{bmatrix} c_{12} \\ c_{22} \\ c_{32} \end{bmatrix} \begin{bmatrix} c_{11}^{-1} & c_{12}^{-1} & c_{13}^{-1} \end{bmatrix}.$$

Therefore we can find non-zero real 1×3 -matrixes $\bar{a}_1, \bar{a}_2$ such that $A = \bar{a}_1^T \bar{a}_2$. As $A^2 = 0$ we have $\bar{a}_1 \perp \bar{a}_2$. Similarly, we can find non-zero real 1×3 -matrixes $\bar{b}_1, \bar{b}_2$ such that $B = \bar{b}_1^T \bar{b}_2$ and $\bar{b}_1 \perp \bar{b}_2$. Let $\bar{o} \in \mathbb{R}^3$ be a non-zero vector that is orthogonal to both $\bar{a}_1$ and $\bar{a}_2$. As $AB = BA = 0$ we have $\bar{a}_1 \perp \bar{b}_2$ and $\bar{a}_2 \perp \bar{b}_1$. It follows that there exists a real matrix $[d_{ij}]_{i,j=1,2}$ such that

$$\begin{aligned} \bar{b}_1 &= d_{11}\bar{a}_1 + d_{12}\bar{o} \\ \bar{b}_2 &= d_{21}\bar{a}_2 + d_{22}\bar{o}. \end{aligned}$$

Then

$$0 = \langle \bar{b}_1, \bar{b}_2 \rangle = d_{12}d_{22}\|\bar{o}\|^2.$$

If $d_{12} = 0$, then $d_{11} \neq 0$ and we can put $\bar{a} := \bar{a}_1, \bar{b} := \bar{a}_2, \bar{c} := d_{11}\bar{b}_2$. Then $\bar{a}^T \bar{b} = A$ and $\bar{a}^T \bar{c} = B$. If $d_{22} = 0$, then $d_{21} \neq 0$ and we can put $\bar{a} := \bar{a}_1/d_{21}, \bar{b} := \bar{b}_1, \bar{c} := \bar{b}_2$. Then $\bar{a}^T \bar{c} = A$ and $\bar{b}^T \bar{c} = B$, which completes the proof. $\square$

By the above lemma, there exists $\bar{c} \in \mathbb{R}^3$ such that for any two linearly independent vectors $\bar{a}, \bar{b} \in \mathbb{R}^3$ orthogonal to $\bar{c}$ there exist C^1 -functions $h_1, h_2 : \mathbb{T}^3 \rightarrow \mathbb{R}$ such that $g(\bar{x})$ equals

$$\bar{c}^T (h_1(\bar{x})\bar{a} + h_2(\bar{x})\bar{b}) \text{ or } (h_1(\bar{x})\bar{a} + h_2(\bar{x})\bar{b})^T \bar{c}$$

for all $\bar{x} \in \mathbb{T}^3$. First we deal with the special case where the limit function g is constant.

Lemma 9 *Let $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ be an area-preserving ergodic C^1 -diffeomorphism with τ -polynomial uniform growth of the derivative ($\tau > 0$). Suppose that the limit function $g = \lim_{n \rightarrow \infty} n^{-\tau} Df^n$ is constant. Then either $\tau = 1$ or $\tau = 2$ and f is algebraically conjugate to a diffeomorphism of the form*

$$\mathbb{T}^3 \ni (x_1, x_2, x_3) \longmapsto (x_1 + \alpha, \varepsilon x_2 + \beta(x_1), x_3 + \gamma(x_1, x_2)) \in \mathbb{T}^3,$$

where $\varepsilon = \det Df = \pm 1$.

Before we will pass to the proof we give notation. Let $A \in GL_3(\mathbb{R})$. Let us denote by $\mathbb{T}_A^3$ the quotient group $\mathbb{R}^3 / \mathbb{Z}^3 A^T$, which is a model of the 3-torus as well. Then the map

$$A : \mathbb{T}^3 \rightarrow \mathbb{T}_A^3, \quad A\bar{x} = \bar{x}A^T$$

establishes a smooth isomorphism between $\mathbb{T}^3$ and $\mathbb{T}_A^3$. Suppose that $\xi : \mathbb{T}_A^3 \rightarrow \mathbb{T}_A^3$ is a diffeomorphism. Then $A^{-1} \circ \xi \circ A$ is a diffeomorphism of the torus $\mathbb{T}^3$. Let $N \in GL_3(\mathbb{Z})$ be its linear part. Then

$$\xi(\bar{x} + \bar{m}A^T) = \xi(\bar{x}) + \bar{m}N^T A^T$$

for all $\bar{m} \in \mathbb{Z}^3$. Moreover, we can decompose

$$\xi(\bar{x}) = \bar{x}(ANA^{-1})^T + \tilde{\xi}(\bar{x})$$

where ANA^{-1} is the linear and $\tilde{\xi}$ is the periodic part of ξ , i.e.

$$\tilde{\xi}(\bar{x} + \bar{m}A^T) = \tilde{\xi}(\bar{x})$$

for all $\bar{m} \in \mathbb{Z}^3$.

Suppose that $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ is a smooth diffeomorphism with τ -polynomial uniform growth of the derivative and $g : \mathbb{T}^3 \rightarrow M_3(\mathbb{R})$ is the limit of the sequence $\{n^{-\tau} Df^n\}_{n \in \mathbb{N}}$. Let us consider the diffeomorphism $\hat{f} : \mathbb{T}_A^3 \rightarrow \mathbb{T}_A^3$ given by $\hat{f} := A \circ f \circ A^{-1}$. Then

$$(11) \quad \frac{1}{n^\tau} D\hat{f}^n(\bar{x}) = \frac{1}{n^\tau} ADf^n(A^{-1}\bar{x})A^{-1} \rightarrow Ag(A^{-1}\bar{x})A^{-1}$$

uniformly on $\mathbb{T}_A^3$. Let us denote by $\hat{g} : \mathbb{T}_A^3 \rightarrow M_3(\mathbb{R})$ the function $\hat{g}(\bar{x}) := Ag(A^{-1}\bar{x})A^{-1}$. Lemma 4 now gives

$$(12) \quad g(\bar{x}) = g(f\bar{x})Df(\bar{x}) \quad \text{and} \quad g(\bar{y}) = Df(\bar{x})g(\bar{y})$$

for all $\bar{x}, \bar{y} \in \mathbb{T}^3$, and consequently

$$(13) \quad \hat{g}(\bar{x}) = \hat{g}(\hat{f}\bar{x})D\hat{f}(\bar{x}) \quad \text{and} \quad \hat{g}(\bar{y}) = D\hat{f}(\bar{x})\hat{g}(\bar{y})$$

for all $\bar{x}, \bar{y} \in \mathbb{T}_A^3$.

Throughout this paper we will denote by $G(\bar{c})$ the subgroup of all $\bar{m} \in \mathbb{Z}^3$ such that $\bar{m} \perp \bar{c}$. Of course, the rank of $G(\bar{c})$ can be equal 0, 1 or 2, whenever $c \neq 0$. Further useful properties of the group $G(\bar{c})$ the reader can find in Appendix B.

Suppose that $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ is an area-preserving ergodic C^1 -diffeomorphism with τ -polynomial uniform growth of the derivative and the limit function g is constant. By Lemma 8, there exist mutually orthogonal vectors $\bar{a}, \bar{c} \in \mathbb{R}^3$ such that $g = \bar{c}^T \bar{a}$. We now prove that following

Lemma 10 *Let $f : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ be an area-preserving C^1 -diffeomorphism. Suppose that f preserves orientation, has τ -polynomial uniform growth of the derivative and the limit function $g = \lim_{n \rightarrow \infty} n^{-\tau} Df^n$ equals $\bar{c}^T \bar{a}$. Then $\text{rank } G(\bar{a}) = 2$. Moreover, τ equals either 1 or 2.*

Proof. Let $\bar{b} \in \mathbb{R}^3$ be a vector orthogonal to both $\bar{a}$ and $\bar{c}$ such that $\det(A) = 1$, where

$$A = \begin{bmatrix} \bar{a} \\ \bar{b} \\ \bar{c} \end{bmatrix}.$$

Consider $\hat{f} : \mathbb{T}_A^3 \rightarrow \mathbb{T}_A^3$ given by $\hat{f} := A \circ f \circ A^{-1}$. Then

$$\hat{g} = A \bar{c}^T \bar{a} A^{-1} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}.$$

From (13) we obtain

$$\begin{bmatrix} 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} D\hat{f} \quad \text{and} \quad \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = D\hat{f} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}.$$

Consequently,

$$\begin{aligned} \frac{\partial}{\partial x_1} \hat{f}_1(\bar{x}) &= 1, & \frac{\partial}{\partial x_2} \hat{f}_1(\bar{x}) &= 0, & \frac{\partial}{\partial x_3} \hat{f}_1(\bar{x}) &= 0, \\ \frac{\partial}{\partial x_3} \hat{f}_1(\bar{x}) &= 0, & \frac{\partial}{\partial x_3} \hat{f}_2(\bar{x}) &= 0, & \frac{\partial}{\partial x_3} \hat{f}_3(\bar{x}) &= 1 \end{aligned}$$

for all $\bar{x} \in \mathbb{T}_A^3$. It follows that

$$\hat{f}(x_1, x_2, x_3) = (x_1 + \alpha, x_2 + \beta(x_1), x_3 + \gamma(x_1, x_2)),$$

where $\beta : \mathbb{R} \rightarrow \mathbb{R}$, $\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$ are C^1 -function. Let $N \in GL_3(\mathbb{Z})$ denote the linear part of $\hat{f}$. Then the linear part of $\hat{f}$ equals

$$ANA^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ K_{21} & 1 & 0 \\ K_{31} & K_{32} & 1 \end{bmatrix}.$$

It follows that

$$(14) \quad \bar{a}N = \bar{a}$$

$$(15) \quad \bar{b}N = K_{21}\bar{a} + \bar{b}$$

$$(16) \quad \bar{c}N = K_{31}\bar{a} + K_{32}\bar{b} + \bar{c}.$$

Let $\tilde{f} : \mathbb{T}^3 \rightarrow \mathbb{R}^3$ denote the periodic part of f , i.e.

$$f(\bar{x}) = \bar{x}N^T + \tilde{f}(\bar{x}).$$

Then

$$f^n(\bar{x}) = \bar{x}(N^n)^T + \sum_{k=0}^{n-1} \tilde{f}(f^k \bar{x})(N^{n-1-k})^T.$$

Since $\int_{\mathbb{T}^3} D(\tilde{f} \circ f^k)(\bar{x})d\bar{x} = 0$ for all natural k , we see that

$$(17) \quad \frac{1}{n^\tau}N^n = \frac{1}{n^\tau} \int_{\mathbb{T}^3} Df^n(\bar{x})d\bar{x} \rightarrow g.$$

It follows that

$$(18) \quad \frac{1}{n^\tau} \begin{bmatrix} 1 & 0 & 0 \\ K_{21} & 1 & 0 \\ K_{31} & K_{32} & 1 \end{bmatrix}^n \rightarrow \hat{g} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

Suppose, contrary to our claim, that $\text{rank } G(\bar{a}) < 2$.

First suppose that $\text{rank } G(\bar{a}) = 0$. From (14) we conclude that $N = \text{Id}$, and finally that $n^{-\tau}N^n$ tends to zero, contrary to (17).

Now suppose that $\text{rank } G(\bar{a}) = 1$. Let $\bar{m} \in \mathbb{Z}^3$ be a generator of $G(\bar{a})$. Then there exists a rational vector $\bar{r} \in \mathbb{Q}^3$ such that $N - \text{Id} = \bar{m}^T \bar{r}$, by (14). From (15) we have

$$\bar{b}\bar{m}^T \bar{r} = \bar{b}(N - \text{Id}) = K_{21}\bar{a}.$$

Suppose that $K_{21} \neq 0$. Then $\text{rank } G(\bar{a}) = \text{rank } G(\bar{r}) = 2$, which contradicts our assumption. Consequently, $K_{21} = 0$. It follows that

$$\begin{bmatrix} 1 & 0 & 0 \\ K_{21} & 1 & 0 \\ K_{31} & K_{32} & 1 \end{bmatrix}^n = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ nK_{31} & nK_{32} & 1 \end{bmatrix}.$$

From (18) it follows that $n = 1$ and $K_{31} = 1, K_{32} = 0$. Then

$$\bar{c}\bar{m}^T \bar{r} = \bar{c}(N - \text{Id}) = \bar{a},$$

by (16). It follows that $\text{rank } G(\bar{a}) = \text{rank } G(\bar{r}) = 2$, which contradicts our assumption.

Finally, we have to prove that τ equals either 1 or 2. From (18) we obtain

$$n^{1-\tau}K_{21} \rightarrow 0, \quad n^{1-\tau}K_{31} + \frac{1-1/n}{2}n^{2-\tau}K_{21}K_{32} \rightarrow 1, \quad n^{1-\tau}K_{32} \rightarrow 0.$$

If $K_{21} = 0$, then $\tau = 1$ and $K_{31} = 1$. Otherwise, $\tau = 2$ and $K_{21}K_{32} = 2$, which completes the proof. $\square$

Proof of Lemma 9. First note that f^2 preserves area and orientation and $n^{-\tau}Df^{2n}$ uniformly tends to $2^\tau \bar{c}^T \bar{a}$. By Lemma 10, $\text{rank } G(\bar{a}) = 2$. It follows that $\bar{a} = a\bar{m} \in a\mathbb{Z}^3$, by Lemma 14 (see Appendix B). Now choose $\bar{n}, \bar{k} \in \mathbb{Z}^3$ such that the determinant of

$$A := \begin{bmatrix} \bar{m} \\ \bar{n} \\ \bar{k} \end{bmatrix}$$

equals 1. Let us consider the diffeomorphism $\hat{f} : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ given by $\hat{f} := A \circ f \circ A^{-1}$. Then

$$\hat{g} = AgA^{-1} = a \begin{bmatrix} 0 \\ \bar{n}\bar{c}^T \\ \bar{k}\bar{c}^T \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}.$$

From (13) we have

$$\begin{bmatrix} 1 & 0 & 0 \end{bmatrix} D\hat{f}(\bar{x}) = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 \\ \bar{n}\bar{c}^T \\ \bar{k}\bar{c}^T \end{bmatrix} = D\hat{f}(\bar{x}) \begin{bmatrix} 0 \\ \bar{n}\bar{c}^T \\ \bar{k}\bar{c}^T \end{bmatrix}.$$

It follows that

$$\hat{f}(x_1, x_2, x_3) = (x_1 + \alpha, \varphi_{x_1}(x_2, x_3)),$$

where $\varphi : \mathbb{T} \times \mathbb{T}^2 \rightarrow \mathbb{T}^2$ is an area-preserving random diffeomorphism over the rotation by an irrational number α . Then

$$\begin{bmatrix} \bar{n}\bar{c}^T \\ \bar{k}\bar{c}^T \end{bmatrix} = D\varphi \begin{bmatrix} \bar{n}\bar{c}^T \\ \bar{k}\bar{c}^T \end{bmatrix}.$$

Suppose that $\bar{n}\bar{c}^T$ and $\bar{k}\bar{c}^T$ are rationally independent. By Lemma 5, $\varphi_{x_1}(x_2, x_3) = (x_2 + \beta(x_1), x_3 + \gamma(x_1))$, where $\beta, \gamma : \mathbb{T} \rightarrow \mathbb{T}$ are C^1 -functions, which is our assertion.

Otherwise, by Lemma 6, there exist a group automorphism $B : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ and C^1 -functions $\beta : \mathbb{T} \rightarrow \mathbb{T}$, $\gamma : \mathbb{T}^2 \rightarrow \mathbb{T}$ such that

$$B \circ \varphi_{x_1} \circ B^{-1}(x_2, x_3) = (\varepsilon x_2 + \beta(x_1), x_3 + \gamma(x_1, x_2)),$$

where $\varepsilon = \det Df$, which implies our assertion. $\square$

Proof of Theorem 7 is divided into a few cases.

Case 1: Suppose that $g = \bar{c}^T(h_1\bar{a} + h_2\bar{b})$, where $\bar{a}$ and $\bar{b}$ are orthogonal to $\bar{c}$ and the matrix

$$A = \begin{bmatrix} \bar{a} \\ \bar{b} \\ \bar{c} \end{bmatrix}$$

is nonsingular. Let $\hat{f} : \mathbb{T}_A^3 \rightarrow \mathbb{T}_A^3$ be given by $\hat{f} := A \circ f \circ A^{-1}$. Then

$$\hat{g} = A\bar{c}^T(\hat{h}_1\bar{a} + \hat{h}_2\bar{b})A^{-1} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \begin{bmatrix} \hat{h}_1 & \hat{h}_2 & 0 \end{bmatrix},$$

where $\hat{h}_i(\bar{x}) := h_i(A^{-1}\bar{x})$ for $i = 1, 2$. From (13) we obtain

$$(19) \quad \begin{bmatrix} \hat{h}_1(\bar{x}) & \hat{h}_2(\bar{x}) & 0 \end{bmatrix} = \begin{bmatrix} \hat{h}_1(\hat{f}\bar{x}) & \hat{h}_2(\hat{f}\bar{x}) & 0 \end{bmatrix} D\hat{f}(\bar{x}),$$

$$\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = D\hat{f}(\bar{x}) \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

for all $\bar{x} \in \mathbb{T}_A^3$. Consequently

$$\frac{\partial}{\partial x_3} \hat{f}_1(\bar{x}) = 0, \quad \frac{\partial}{\partial x_3} \hat{f}_2(\bar{x}) = 0 \quad \text{and} \quad \frac{\partial}{\partial x_3} \hat{f}_3(\bar{x}) = 1$$

for all $\bar{x} \in \mathbb{T}_A^3$. It follows that

$$\hat{f}(x_1, x_2, x_3) = (F(x_1, x_2), x_3 + \gamma(x_1, x_2)),$$

where $\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a smooth function and $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the diffeomorphism given by $F(x_1, x_2) = (\hat{f}_1(x_1, x_2), \hat{f}_2(x_1, x_2))$. Let K denote the linear part of $\hat{f}$. Then $K = A N A^{-1}$, where $N \in GL_3(\mathbb{Z})$ is the linear part of f . Then $\det K = \det N = \varepsilon' = \pm 1$ and $K_{13} = 0, K_{23} = 0, K_{33} = 1$. Moreover, there exist $(\bar{a}\bar{m}^T, \bar{b}\bar{m}^T)$ -periodic for all $\bar{m} \in \mathbb{Z}^3$ functions $\tilde{F} : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \tilde{\gamma} : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that

$$\begin{aligned} F(\bar{x}) &= \tilde{F}(\bar{x}) + \bar{x}K'^T, \\ \gamma(x_1, x_2) &= \tilde{\gamma}(x_1, x_2) + K_{31}x_1 + K_{32}x_2, \end{aligned}$$

where $K' = K|_{\{1,2\} \times \{1,2\}} \in GL_2(\mathbb{R})$ and $\det K' = \varepsilon'$. From (11) we deduce that

$$\frac{1}{n^\tau} DF^n(x_1, x_2) \rightarrow 0 \quad \text{and} \quad \frac{1}{n^\tau} \sum_{k=0}^{n-1} D(\gamma \circ F^k)(x_1, x_2) \rightarrow \begin{bmatrix} \hat{h}_1(\bar{x}) & \hat{h}_2(\bar{x}) \end{bmatrix}$$

uniformly on $\mathbb{T}_A^3$. Therefore $\hat{h}_1, \hat{h}_2$ depend only on the first two coordinates. Let $H : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be given by $H(x_1, x_2) = \begin{bmatrix} \hat{h}_1(x_1, x_2, 0) & \hat{h}_2(x_1, x_2, 0) \end{bmatrix}$. Then H is $(\bar{a}\bar{m}^T, \bar{b}\bar{m}^T)$ -periodic for all $\bar{m} \in \mathbb{Z}^3$. From (19) we have

$$(20) \quad H(F\bar{x})DF(\bar{x}) = H(\bar{x})$$

for all $\bar{x} \in \mathbb{R}^2$. Set $\chi_n := n^{-\tau} \sum_{k=0}^{n-1} \gamma \circ F^k$. Since $D\chi_n \rightarrow H$ uniformly on $\mathbb{R}^2$, we see that

$$\begin{aligned} \chi_n(x_1, x_2) - \chi_n(x_1, 0) &\rightarrow \int_0^{x_2} H_2(x_1, t) dt, \\ \chi_n(x_1, x_2) - \chi_n(0, x_2) &\rightarrow \int_0^{x_1} H_1(t, x_2) dt \end{aligned}$$

for all $(x_1, x_2) \in \mathbb{R}^2$. Let $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$\xi(x_1, x_2) := \lim_{n \rightarrow \infty} (\chi_n(x_1, x_2) - \chi_n(0, 0)) = \begin{cases} \int_0^{x_1} H_1(t, x_2) dt + \int_0^{x_2} H_2(0, t) dt \\ \int_0^{x_2} H_2(x_1, t) dt + \int_0^{x_1} H_1(t, 0) dt. \end{cases}$$

Then $\partial\xi/\partial x_1 = H_1$, $\partial\xi/\partial x_2 = H_2$ and ξ is of class C^2 . From (20) it follows that there exists $\alpha \in \mathbb{R}$ such that

$$(21) \quad \xi(F\bar{x}) = \xi(\bar{x}) + \alpha.$$

By Lemma 14 (see Appendix B), there exist a $(\bar{a}\bar{m}^T, \bar{b}\bar{m}^T)$ -periodic for all $\bar{m} \in \mathbb{Z}^3$ function $\tilde{\xi} : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $d_1, d_2 \in \mathbb{R}$ such that $\xi(x_1, x_2) = \tilde{\xi}(x_1, x_2) + d_1x_1 + d_2x_2$. Since $H \neq 0$, it is easy to see that $(d_1, d_2) \neq (0, 0)$. Moreover, from (21) we have

$$(22) \quad \begin{bmatrix} d_1 & d_2 \end{bmatrix} K' = \begin{bmatrix} d_1 & d_2 \end{bmatrix}$$

and

$$(23) \quad \begin{aligned} & \tilde{\xi}(\bar{x}) + \alpha \\ &= \tilde{\xi}(\tilde{F}_1(\bar{x}) + K_{11}x_1 + K_{12}x_2, \tilde{F}_2(\bar{x}) + K_{21}x_1 + K_{22}x_2) + d_1\tilde{F}_1(\bar{x}) + d_2\tilde{F}_2(\bar{x}). \end{aligned}$$

Case 1a: Suppose that $\text{rank } G(\bar{c}) = 0$. By Lemma 14, $D\hat{f}$ is constant. It follows that Df and g are constant. Therefore $g = \bar{c}^T \bar{a}$, where $\bar{a}$ is orthogonal to $\bar{c}$. From (12) we obtain $\bar{c}^T = Df \bar{c}^T$. As $G(\bar{c}) = \{0\}$ we have $Df = \text{Id}$. Consequently, f is a rotation on the 3-torus, which is impossible.

Case 1b: Suppose that $\text{rank } G(\bar{c}) = 1$. By Lemma 14, there exist $\bar{F} : \mathbb{T} \rightarrow \mathbb{R}^2$, $\bar{\xi} : \mathbb{T} \rightarrow \mathbb{R}$, $\bar{\gamma} : \mathbb{T} \rightarrow \mathbb{R}$ and real numbers l_1, l_2 such that $\bar{m} = l_1\bar{a} + l_2\bar{b}$ generates $G(\bar{c})$ and

$$\begin{aligned} \tilde{F}(x_1, x_2) &= \bar{F}(l_1x_1 + l_2x_2), \\ \tilde{\xi}(x_1, x_2) &= \bar{\xi}(l_1x_1 + l_2x_2), \\ \tilde{\gamma}(x_1, x_2) &= \bar{\gamma}(l_1x_1 + l_2x_2). \end{aligned}$$

From (23) we obtain

$$\begin{aligned} & \bar{\xi}(l_1x_1 + l_2x_2) + \alpha \\ &= \bar{\xi}(l_1\bar{F}_1(l_1x_1 + l_2x_2) + l_2\bar{F}_2(l_1x_1 + l_2x_2) + s_1x_1 + s_2x_2) \\ & \quad + d_1\bar{F}_1(l_1x_1 + l_2x_2) + d_2\bar{F}_2(l_1x_1 + l_2x_2), \end{aligned}$$

where $\begin{bmatrix} s_1 & s_2 \end{bmatrix} = \begin{bmatrix} l_1 & l_2 \end{bmatrix} K'$. If (s_1, s_2) and (l_1, l_2) are linearly independent, then $\bar{\xi}$ is constant. It follows that H is constant and we can come back to Lemma 9. Otherwise, there exists a real number s such that $(s_1, s_2) = s(l_1, l_2)$ and

$$\bar{\xi}(x) + \alpha = \bar{\xi}(l_1\bar{F}_1(x) + l_2\bar{F}_2(x) + sx) + d_1\bar{F}_1(x) + d_2\bar{F}_2(x)$$

for any real x . As f preserves area we have $\det DF(\bar{x}) = \varepsilon = \pm 1$. It follows that

$$\begin{aligned}\varepsilon &= \det \begin{bmatrix} l_1 D\bar{F}_1(x) + K_{11} & l_2 D\bar{F}_1(x) + K_{12} \\ l_1 D\bar{F}_2(x) + K_{21} & l_2 D\bar{F}_2(x) + K_{22} \end{bmatrix} \\ &= (l_1 K_{22} - l_2 K_{21}) D\bar{F}_1(x) + (-l_1 K_{12} + l_2 K_{11}) D\bar{F}_2(x) + \det K \\ &= (l_1 D\bar{F}_1(x) + l_2 D\bar{F}_2(x)) \det K/s + \det K\end{aligned}$$

for any real x . Since $\bar{F}_1, \bar{F}_2$ are 1-periodic, we have $l_1 D\bar{F}_1(x) + l_2 D\bar{F}_2(x) = 0$ and $\det K = \varepsilon$. Therefore the function $l_1 \bar{F}_1 + l_2 \bar{F}_2$ is constant. Let us choose real numbers r_1, r_2 such that the determinant of the matrix

$$L = \begin{bmatrix} l_1 & l_2 & 0 \\ r_1 & r_2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

equals 1. Now consider the diffeomorphism $\check{f} : \mathbb{T}_{LA}^3 \rightarrow \mathbb{T}_{LA}^3$ given by $\check{f} = L \circ \hat{f} \circ L^{-1}$. Then

$$\check{f}(x_1, x_2, x_3) = (sx_1 + \alpha, \varepsilon/sx_2 + rx_1 + r_1\bar{F}_1(x_1) + r_2\bar{F}_2(x_1), x_3 + \bar{\gamma}(x_1) + p_1x_1 + p_2x_2).$$

As $\partial \check{f}_1^n / \partial x_1 = s^n$ and $\partial \check{f}_2^n / \partial x_2 = (\varepsilon/s)^n$ we obtain $s = \pm 1$, because $\check{f}$ has polynomial uniform growth of the derivative. Moreover,

$$LA = \begin{bmatrix} \bar{m} \\ r_1\bar{a} + r_2\bar{b} \\ \bar{c} \end{bmatrix}$$

and $LA \circ f = \check{f} \circ LA$. Therefore $f(\bar{x})\bar{m}^T = s\bar{x}\bar{m}^T + \alpha$. Observe that $s = 1$. Indeed, suppose, contrary to our claim, that $s = -1$. Consider the smooth function $\kappa : \mathbb{T}^3 \rightarrow \mathbb{C}$ given by $\kappa(\bar{x}) = e^{2\pi i \bar{x}\bar{m}^T}$. Then $\kappa \circ f^2 = \kappa$. Since κ is smooth, we conclude that is constant, by the ergodicity of f . Consequently, $\bar{m} = 0$, which is impossible.

Now choose $\bar{n}, \bar{k} \in \mathbb{Z}^3$ such that the determinant of

$$A := \begin{bmatrix} \bar{m} \\ \bar{n} \\ \bar{k} \end{bmatrix}$$

equals 1. Let us consider the diffeomorphism $\hat{f} : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ given by $\hat{f} := A \circ f \circ A^{-1}$. From (13) we have

$$\begin{bmatrix} 0 \\ \bar{n}\bar{c}^T \\ \bar{k}\bar{c}^T \end{bmatrix} = D\hat{f}(\bar{x}) \begin{bmatrix} 0 \\ \bar{n}\bar{c}^T \\ \bar{k}\bar{c}^T \end{bmatrix}.$$

Moreover,

$$\hat{f}_1(\bar{x}) = f(\bar{x}(A^{-1})^T)\bar{m}^T = \bar{x}(A^{-1})^T\bar{m}^T + \alpha = x_1 + \alpha.$$

Now our claim follows by the same method as in the proof of Lemma 9.

Case 1c: Suppose that $\text{rank } G(\bar{c}) = 2$. Then we can assume that $\bar{a}, \bar{b}, \bar{c} \in \mathbb{Z}^3$ and $\bar{a}, \bar{b}$ generates $G(\bar{c})$. Set $q = \det A \in \mathbb{N}$. Then the linear part of $\hat{f}$ (equals $K = A N A^{-1}$) belongs to $M_3(\mathbb{Z}/q)$. Moreover, the functions $\tilde{F} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $\tilde{\gamma} : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $\tilde{\xi} : \mathbb{R}^2 \rightarrow \mathbb{R}$ are $\mathbb{Z}^2$ -periodic, by Corollary 1 (see Appendix B).

Case 1c(i): Suppose that d_1/d_2 is irrational. From (22) we obtain $K' = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$. Set

$$L := \begin{bmatrix} 1/q & 0 & 0 \\ 0 & 1/q & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Now consider the diffeomorphism $\check{f} : \mathbb{T}_{LA}^3 \rightarrow \mathbb{T}_{LA}^3$ given by $\check{f} = L \circ \hat{f} \circ L^{-1}$. Then

$$\check{f}(x_1, x_2, x_3) = (\tilde{F}(x_1, x_2), x_3 + \tilde{\gamma}(x_1, x_2)),$$

where

$$\begin{aligned} \tilde{F}(x_1, x_2) &= \frac{1}{q} F(qx_1, qx_2), \\ \tilde{\gamma}(x_1, x_2) &= \gamma(qx_1, qx_2). \end{aligned}$$

Then

$$\begin{aligned} \tilde{F}(\bar{x} + \bar{m}) - \tilde{F}(\bar{x}) &= \bar{m}, \\ \tilde{\gamma}(\bar{x} + \bar{m}) - \tilde{\gamma}(\bar{x}) &= qK_{31}m_1 + qK_{32}m_2 \in \mathbb{Z} \end{aligned}$$

for all $\bar{m} \in \mathbb{Z}^2$. Therefore, $\check{f}$ can be also treated as a diffeomorphism of the torus $\mathbb{T}^3$. Let $\check{\xi}(x_1, x_2) = \xi(qx_1, qx_2)$. Then

$$(24) \quad \check{\xi} \circ \check{F} = \check{\xi} + \alpha,$$

$D\check{\xi} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is $\mathbb{Z}^2$ -periodic and non-zero at each point. Moreover, $\check{f} : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ has τ -polynomial uniform growth of the derivative. Precisely,

$$(25) \quad \frac{1}{n^\tau} D\check{f}^n \rightarrow \begin{bmatrix} 0 & 0 \\ D\check{\xi} & 0 \end{bmatrix}$$

uniformly.

Let us denote by φ^t the Hamiltonian C^2 -flow on $\mathbb{T}^2$ defined by the Hamiltonian equation

$$\frac{d}{dt} \varphi^t(\bar{x}) = \begin{bmatrix} \check{\xi}_{x_2}(\varphi^t(\bar{x})) \\ -\check{\xi}_{x_1}(\varphi^t(\bar{x})) \end{bmatrix}.$$

Since φ^t has no fixed point and

$$\int_{\mathbb{T}^2} \check{\xi}_{x_1}(\bar{x}) d\bar{x} / \int_{\mathbb{T}^2} \check{\xi}_{x_2}(\bar{x}) d\bar{x} = d_1/d_2$$

is irrational, it follows that φ^t is C^2 -conjugate to the special flow constructed over the rotation by an irrational number a and under a positive C^2 -function $b : \mathbb{T} \rightarrow \mathbb{R}$ with $\int_{\mathbb{T}} b(x) dx = 1$, (see for instance [2, Ch. 16]) i.e. there exists an area-preserving C^2 -diffeomorphism $\rho : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and a matrix $N \in GL_2(\mathbb{Z})$ such that

$$\sigma^t \circ \rho = \rho \circ \varphi^t,$$

where $\sigma^t(x_1, x_2) = (x_1, x_2 + t)$ and

$$\rho(\bar{x} + \bar{m}) = (\rho_1(\bar{x}) + (\bar{m}N)_1 + (\bar{m}N)_2 a, \rho_2(\bar{x}) - b^{((\bar{m}N)_2)}(\rho_1(\bar{x})))$$

for all $\bar{m} \in \mathbb{Z}^2$. Let $T_{a,-b} : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$ denote the skew product given by $T_{a,-b}(x_1, x_2) = (x_1 + a, x_2 - b(x_1))$. Let us consider the quotient space $M = \mathbb{T} \times \mathbb{R} / \sim$, where the relation $\sim$ is defined by $(x_1, x_2) \sim (y_1, y_2)$ iff $(x_1, x_2) = T_{a,-b}^k(y_1, y_2)$ for an integer k . Then the quotient flow of the action σ^t by the relation $\sim$ is the special flow constructed over the rotation by a and under the function b . Moreover, $\rho : \mathbb{T}^2 \rightarrow M$ conjugates flows φ^t and σ^t . Let $\bar{F} : M \rightarrow M$ denote the C^2 -diffeomorphism given by $\bar{F} := \rho \circ \tilde{F} \circ \rho^{-1}$. Since the map $\mathbb{R} \ni t \mapsto \tilde{\xi}(\varphi^t \bar{x}) \in \mathbb{R}$ is constant for each $\bar{x} \in \mathbb{R}^2$ we see that the map

$$\mathbb{R} \ni t \mapsto \tilde{\xi} \circ \rho^{-1}(\sigma^t(x_1, x_2)) = \tilde{\xi} \circ \rho^{-1}(x_1, x_2 + t) \in \mathbb{R}$$

is constant for each $(x_1, x_2) \in \mathbb{R}^2$. It follows that the function $\tilde{\xi} \circ \rho^{-1} : \mathbb{R}^2 \rightarrow \mathbb{R}$ depends only on the first coordinate. Moreover,

$$\begin{aligned} D\rho^{-1}(\bar{x}) \begin{bmatrix} 0 \\ 1 \end{bmatrix} &= \frac{d}{dt} \rho^{-1} \circ \sigma^t(\bar{x})|_{t=0} = \frac{d}{dt} \varphi^t \circ \rho^{-1}(\bar{x})|_{t=0} \\ &= \begin{bmatrix} \tilde{\xi}_{x_2}(\rho^{-1}(\bar{x})) \\ -\tilde{\xi}_{x_1}(\rho^{-1}(\bar{x})) \end{bmatrix}. \end{aligned}$$

Consequently, $\partial \rho_1^{-1} / \partial x_2 = \partial \tilde{\xi} / \partial x_2 \circ \rho^{-1}$ and $\partial \rho_2^{-1} / \partial x_2 = -\partial \tilde{\xi} / \partial x_1 \circ \rho^{-1}$. It follows that

$$\frac{d}{dx_1} \tilde{\xi} \circ \rho^{-1} = \frac{\partial \tilde{\xi}}{\partial x_1} \circ \rho^{-1} \cdot \frac{\partial \rho_1^{-1}}{\partial x_1} + \frac{\partial \tilde{\xi}}{\partial x_2} \circ \rho^{-1} \cdot \frac{\partial \rho_2^{-1}}{\partial x_1} = -\det D\rho^{-1} = \delta = \pm 1.$$

Therefore

$$(26) \quad \tilde{\xi} \circ \rho^{-1}(x_1, x_2) = \delta x_1 + c.$$

From (24) we see that $\tilde{\xi} \circ \rho^{-1} \circ \bar{F} = \tilde{\xi} \circ \rho^{-1} + \alpha$ and consequently that $\bar{F}_1(x_1, x_2) = x_1 + \delta \alpha$. To shorten notation, we will write α instead of $\delta \alpha$. Since $\bar{F} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ preserves area, we conclude that

$$\bar{F}(x_1, x_2) = (x_1 + \alpha, \varepsilon x_2 + \beta(x_1)),$$

where $\beta : \mathbb{R} \rightarrow \mathbb{R}$ is a C^2 -function and $\varepsilon = \det D\bar{F} = \pm 1$. As $\bar{F}$ is a diffeomorphism of M , there exists $m_1, m_2 \in \mathbb{Z}$ such that

$$\begin{aligned} (x_1 + 1 + \alpha, \varepsilon x_2 + \beta(x_1 + 1)) &= \bar{F}(x_1 + 1, x_2) = T_{a,-b}^{m_2} \bar{F}(x_1, x_2) + (m_1, 0) \\ &= (x_1 + \alpha + m_1 + m_2 a, \varepsilon x_2 + \beta(x_1) - b^{(m_2)}(x_1 + \alpha)). \end{aligned}$$

It follows that $m_1 = 1$, $m_2 = 0$, hence that $\beta : \mathbb{T} \rightarrow \mathbb{R}$. Moreover, there exists $n_1, n_2 \in \mathbb{Z}$ such that

$$\begin{aligned} & (x_1 + a + \alpha, \varepsilon x_2 - \varepsilon b(x_1) + \beta(x_1 + a)) \\ &= \bar{F} \circ T_{a,-b}(x_1, x_2) = T_{a,-b}^{n_2} \bar{F}(x_1, x_2) + (n_1, 0) \\ &= (x_1 + \alpha + n_1 + n_2 a, \varepsilon x_2 + \beta(x_1) - b^{(n_2)}(x_1 + \alpha)). \end{aligned}$$

It follows that $n_1 = 0$, $n_2 = 1$, hence that $\beta(x) - b(x + \alpha) = -\varepsilon b(x) + \beta(x + a)$. Hence

$$1 - \varepsilon = \int_{\mathbb{T}} (b(x + \alpha) - \varepsilon b(x)) dx = \int_{\mathbb{T}} (\beta(x) - \beta(x + a)) dx = 0.$$

Therefore

$$\bar{F}(x_1, x_2) = (x_1 + \alpha, x_2 + \beta(x_1))$$

and the skew products $\bar{F}$ and $T_{a,-b}$ commute.

Let us consider the diffeomorphism $\bar{f} : M \times \mathbb{T} \rightarrow M \times \mathbb{T}$ defined by

$$\bar{f} := \rho \times \text{Id}_{\mathbb{T}} \circ \check{f} \circ (\rho \times \text{Id}_{\mathbb{T}})^{-1}.$$

Then

$$\bar{f}(x_1, x_2, x_3) = (\bar{F}(x_1, x_2), x_3 + \bar{\gamma}(x_1, x_2)),$$

where $\bar{\gamma} : M \rightarrow \mathbb{T}$ is given by $\bar{\gamma} = \check{\gamma} \circ \rho^{-1}$. Therefore there exist $k_1, k_2 \in \mathbb{Z}$ such that

$$\begin{aligned} \bar{\gamma}(x_1 + 1, x_2) &= \bar{\gamma}(x_1, x_2) + k_1 \\ \bar{\gamma}(x_1 + a, x_2 - b(x_1)) &= \bar{\gamma}(x_1, x_2) + k_2. \end{aligned}$$

Moreover,

$$\begin{aligned} \frac{1}{n^\tau} D \bar{f}^n &= \begin{bmatrix} D\rho \circ \check{F}^n \circ \rho^{-1} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} n^{-\tau} D \check{F}^n \circ \rho^{-1} & 0 \\ n^{-\tau} D(\check{\gamma}^{(n)}) \circ \rho^{-1} & n^{-\tau} \end{bmatrix} \begin{bmatrix} D\rho^{-1} & 0 \\ 0 & 1 \end{bmatrix} \\ &\rightarrow \begin{bmatrix} 0 & 0 \\ D\check{\xi} \circ \rho^{-1} & 0 \end{bmatrix} \begin{bmatrix} D\rho^{-1} & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ D(\check{\xi} \circ \rho^{-1}) & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \delta & 0 & 0 \end{bmatrix} \end{aligned}$$

uniformly, by (25) and (26). It follows that

$$\frac{1}{n^\tau} \sum_{k=0}^{n-1} (\bar{\gamma}_{x_1} \circ \bar{F}^k + \bar{\gamma}_{x_2} \circ \bar{F}^k \cdot D\beta^{(k)}) \rightarrow \delta$$

and

$$\frac{1}{n^\tau} \sum_{k=0}^{n-1} \bar{\gamma}_{x_2} \circ \bar{F}^k \rightarrow 0$$

uniformly, hence that

$$\frac{1}{n^\tau} \sum_{k=0}^{n-1} \int_M (\bar{\gamma}_{x_1}(\bar{F}^k(x_1, x_2)) + \bar{\gamma}_{x_2}(\bar{F}^k(x_1, x_2))) D\beta^{(k)}(x_1) dx_1 dx_2 \rightarrow \delta$$

and

$$(27) \quad \frac{1}{n^\tau} \sum_{k=0}^{n-1} \int_M \bar{\gamma}_{x_2}(\bar{F}^k(x_1, x_2)) dx_1 dx_2 \rightarrow 0.$$

We show that

$$\frac{1}{n} \sum_{k=0}^{n-1} \int_M (\bar{\gamma}_{x_1}(\bar{F}^k(x_1, x_2)) + \bar{\gamma}_{x_2}(\bar{F}^k(x_1, x_2))) D\beta^{(k)}(x_1) dx_1 dx_2 \rightarrow b(0)k_1.$$

This gives $\tau = 1$ and $k_1 \neq 0$. To prove it, note that

$$\begin{aligned} & \frac{1}{n} \sum_{k=0}^{n-1} \int_M \bar{\gamma}_{x_1}(\bar{F}^k(x_1, x_2)) dx_1 dx_2 \\ &= \int_0^1 \int_0^{b(x_1)} \bar{\gamma}_{x_1}(x_1, x_2) dx_2 dx_1 \\ &= \int_0^1 \frac{d}{dx_1} \int_0^{b(x_1)} \bar{\gamma}(x_1, x_2) dx_2 dx_1 - \int_0^1 Db(x_1) \bar{\gamma}(x_1, b(x_1)) dx_1 \\ &= \int_0^{b(1)} \bar{\gamma}(1, x_2) dx_2 - \int_0^{b(0)} \bar{\gamma}(0, x_2) dx_2 - \int_0^1 Db(x_1) (\bar{\gamma}(x_1 + a, 0) - k_2) dx_1 \\ &= b(0)k_1 - \int_0^1 Db(x_1) \bar{\gamma}(x_1 + a, 0) dx_1. \end{aligned}$$

Next observe that

$$\begin{aligned} & \frac{1}{n} \sum_{k=0}^{n-1} \int_M \bar{\gamma}_{x_2}(\bar{F}^k(x_1, x_2)) D\beta^{(k)}(x_1) dx_1 dx_2 \\ &= \frac{1}{n} \sum_{k=0}^{n-1} \int_0^1 \int_0^{b(x_1)} \bar{\gamma}_{x_2}(x_1, x_2) D\beta^{(k)}(x_1 - k\alpha) dx_2 dx_1 \\ &= \frac{1}{n} \sum_{k=0}^{n-1} \int_0^1 (\bar{\gamma}(x_1, b(x_1)) - \bar{\gamma}(x_1, 0)) D\beta^{(k)}(x_1 - k\alpha) dx_1 \\ &= \frac{1}{n} \sum_{k=0}^{n-1} \int_0^1 (\bar{\gamma}(x_1 + a, 0) - k_2 - \bar{\gamma}(x_1, 0)) D\beta^{(k)}(x_1 - k\alpha) dx_1 \\ &= \frac{1}{n} \sum_{k=0}^{n-1} \int_0^1 \bar{\gamma}(x_1 + a, 0) (D\beta^{(k)}(x_1 - k\alpha) - D\beta^{(k)}(x_1 - k\alpha + a)) dx_1 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{n} \sum_{k=0}^{n-1} \int_0^1 \bar{\gamma}(x_1 + a, 0) (Db(x_1) - Db(x_1 - k\alpha)) dx_1 \\
&= \int_0^1 \bar{\gamma}(x_1 + a, 0) Db(x_1) dx_1 - \int_0^1 \bar{\gamma}(x_1 + a, 0) \frac{1}{n} \sum_{k=0}^{n-1} Db(x_1 - k\alpha) dx_1 \\
&\rightarrow \int_0^1 \bar{\gamma}(x_1 + a, 0) Db(x_1) dx_1,
\end{aligned}$$

which proves the desire conclusion. From (27) we have

$$\int_0^1 \int_0^{b(x_1)} \bar{\gamma}_{x_2}(x_1, x_2) dx_2 dx_1 = 0.$$

However

$$\begin{aligned}
\int_0^1 \int_0^{b(x_1)} \bar{\gamma}_{x_2}(x_1, x_2) dx_2 dx_1 &= \int_0^1 (\bar{\gamma}(x_1, b(x_1)) - \bar{\gamma}(x_1, 0)) dx_1 \\
&= \int_0^1 (\bar{\gamma}(x_1 + a, 0) - \bar{\gamma}(x_1, 0) - k_2) dx_1 \\
&= k_1 a - k_2.
\end{aligned}$$

It follows that $k_1 a = k_2$, which contradicts the fact that $k_1 \neq 0$ and a is irrational. Consequently, d_1/d_2 must be rational.

Case 1c(ii): Suppose that $(d_1, d_2) = d(l_1, l_2)$, where l_1, l_2 are relatively prime integer numbers. Then there exist $M \in GL_2(\mathbb{Z})$ and $m \in \mathbb{Z}$ such that

$$K' = M^{-1} \begin{bmatrix} 1 & 0 \\ m/q & \varepsilon \end{bmatrix} M,$$

by (22). Hence there exists an even number r such that $K'^r \in GL_2(\mathbb{Z})$. Therefore the diffeomorphism $F^r : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ can be treated as an area-preserving diffeomorphism of the torus $\mathbb{T}^2$. Let $\check{\xi} : \mathbb{T}^2 \rightarrow \mathbb{T}$ be given by $\check{\xi}(x_1, x_2) = d^{-1}\xi(x_1, x_2)$. Then

$$\check{\xi} \circ F^r = \check{\xi} + r\alpha/d \quad \text{and} \quad (d_1(\check{\xi}), d_2(\check{\xi})) = (l_1, l_2) \neq 0.$$

Notice that α/d is irrational. Indeed, suppose that $\alpha/d = k/l$, where $k \in \mathbb{Z}$ and $l \in \mathbb{N}$. Let $\Xi : \mathbb{T}_A^3 \rightarrow \mathbb{C}$ be given by

$$\Xi(x_1, x_2, x_3) = \exp 2\pi i l \check{\xi}(x_1, x_2).$$

As $\check{\xi} \circ F = \check{\xi} + k/l$, we have

$$\Xi \hat{f}(x_1, x_2, x_3) = \exp 2\pi i l \check{\xi} \circ F(x_1, x_2) = \Xi(x_1, x_2, x_3).$$

By the ergodicity of $\hat{f}$, Ξ and finally $\check{\xi}$ is constant, which is impossible.

By Theorem 13 (see Appendix A), there exists an area-preserving C^2 -diffeomorphism $\psi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ such that

$$\psi^{-1} \circ F^r \circ \psi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$$

is a skew product and $\check{\xi} \circ \psi(x_1, x_2) = x_1$. Therefore $D(\xi \circ \psi) = [d \ 0]$. Let $L \in GL_2(\mathbb{Z})$ denote the linear part of ψ . Set

$$\bar{L} := \begin{bmatrix} L & 0 \\ 0 & 1 \end{bmatrix} \in GL_3(\mathbb{Z}).$$

Let us consider the area-preserving C^2 -isomorphism $\rho : \mathbb{T}_A^3 \rightarrow \mathbb{T}_{\bar{L}^{-1}A}^3$ defined by

$$\rho(x_1, x_2, x_3) = (\psi^{-1}(x_1, x_2), x_3).$$

Let $\check{f} : \mathbb{T}_{\bar{L}^{-1}A}^3 \rightarrow \mathbb{T}_{\bar{L}^{-1}A}^3$ be given by $\check{f} = \rho \circ \hat{f} \circ \rho^{-1}$. Then

$$\begin{aligned} \frac{1}{n^\tau} D\check{f}^n &= \begin{bmatrix} D\psi^{-1} \circ F^n \circ \psi & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} n^{-\tau} DF^n \circ \psi & 0 \\ n^{-\tau} D(\gamma^{(n)}) \circ \psi & n^{-\tau} \end{bmatrix} \begin{bmatrix} D\psi & 0 \\ 0 & 1 \end{bmatrix} \\ &\rightarrow \begin{bmatrix} 0 & 0 \\ D\xi \circ \psi & 0 \end{bmatrix} \begin{bmatrix} D\psi & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ D(\xi \circ \psi) & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ d & 0 & 0 \end{bmatrix} \end{aligned}$$

uniformly. Let us consider the diffeomorphism $\bar{f} : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ given by $\bar{f} = A^{-1} \bar{L} \circ \check{f} \circ \bar{L}^{-1} A$. It is easy to see that

$$\frac{1}{n^\tau} D\bar{f}^n \rightarrow A^{-1} \bar{L} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ d & 0 & 0 \end{bmatrix} \bar{L}^{-1} A$$

uniformly and that $\bar{f}$ and f are conjugate via the area-preserving C^2 -diffeomorphism $A^{-1} \bar{L} \circ \rho \circ A : \mathbb{T}^3 \rightarrow \mathbb{T}^3$. Now applying Lemma 9 for $\bar{f}$ gives our claim.

Case 2: Suppose that $g = (h_1 \bar{a}^T + h_2 \bar{b}^T) \bar{c}$, where $\bar{a}$ and $\bar{b}$ are orthogonal to $\bar{c}$ and the determinant of the matrix

$$A^{-1} = \begin{bmatrix} \bar{c}^T & \bar{a}^T & \bar{b}^T \end{bmatrix}$$

equals 1. Let $\hat{f} : \mathbb{T}_A^3 \rightarrow \mathbb{T}_A^3$ is given by $\hat{f} := A \circ f \circ A^{-1}$. Then

$$\hat{g} = A(h_1 \bar{a}^T + h_2 \bar{b}^T) \bar{c} A^{-1} = \begin{bmatrix} 0 \\ \hat{h}_1 \\ \hat{h}_2 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \end{bmatrix},$$

where $\hat{h}_i(\bar{x}) := h_i(A^{-1} \bar{x})$ for $i = 1, 2$. From (13) we obtain

$$\begin{bmatrix} 0 \\ \hat{h}_1(\bar{x}) \\ \hat{h}_2(\bar{x}) \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \hat{h}_1(\hat{f} \bar{x}) \\ \hat{h}_2(\hat{f} \bar{x}) \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} D\hat{f}(\bar{x})$$

for all $\bar{x} \in \mathbb{T}_A^3$. Consequently

$$\frac{\partial}{\partial x_1} \hat{f}_1(\bar{x}) \hat{h}_i(\hat{f}\bar{x}) = \hat{h}_i(\bar{x}), \quad \frac{\partial}{\partial x_2} \hat{f}_1(\bar{x}) = \frac{\partial}{\partial x_3} \hat{f}_1(\bar{x}) = 0 \quad \text{and} \quad \frac{\partial}{\partial x_1} \hat{f}_1(\bar{x}) \neq 0$$

for all $\bar{x} \in \mathbb{T}_A^3$ and $i = 1, 2$. Now observe that $\hat{h}_1, \hat{h}_2$ are linearly dependent. Indeed, without loss of generality we can assume that $\hat{h}_2$ is $A\lambda^{\otimes 3}$ non-zero. Then $\hat{h}_2(\bar{x}) \neq 0$ for a.e. $\bar{x} \in \mathbb{T}_A^3$, by the ergodicity of $\hat{f}$. Therefore the measurable function $\hat{h}_1/\hat{h}_2 : \mathbb{T}_A^3 \rightarrow \mathbb{R}$ is $\hat{f}$ -invariant. Hence there is a real constant c such that $\hat{h}_1(\bar{x}) = c\hat{h}_2(\bar{x})$ for a.e. $\bar{x} \in \mathbb{T}_A^3$, by ergodicity. Consequently, $h_1 = ch_2$, which leads us to Case 1, and the proof is complete. $\square$

4 4-dimensional case

In this section we indicate why there is no 4-dimensional analogue of classifications of area-preserving diffeomorphisms of polynomial growth of the derivative presented in previous sections. Precisely, we construct an ergodic area-preserving diffeomorphism of the 4-dimensional torus with linear uniform growth of the derivative which is not even metrically isomorphic to any 3-steps skew product, i.e. to any automorphism of $\mathbb{T}^4$ of the form

$$\mathbb{T}^4 \ni (x_1, x_2, x_3, x_4) \longmapsto (x_1 + \alpha, \varepsilon_1 x_2 + \beta(x_1), \varepsilon_2 x_3 + \gamma(x_1, x_2), \varepsilon_3 x_4 + \delta(x_1, x_2, x_3)) \in \mathbb{T}^4,$$

where $\varepsilon_i = \pm 1$ for $i = 1, 2, 3$. Before we pass to the 4-dimensional case we should mention area preserving diffeomorphisms of the 2-torus with sublinear growth of the derivative.

Definition 2 We say that a C^1 -diffeomorphism $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ has *sublinear growth of the derivative* if the sequence Df^n/n tends uniformly to zero.

Suppose that $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ is an area-preserving weakly mixing C^∞ -diffeomorphism with sublinear growth of the derivative. The examples of such diffeomorphisms will be given later. Let $T_\varphi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be an Anzai skew product where $Tx = x + \alpha$ is an ergodic rotation on the circle and $\varphi : \mathbb{T} \rightarrow \mathbb{T}$ is a C^∞ -function with non-zero topological degree.

Theorem 11 *The product diffeomorphism $f \times T_\varphi : \mathbb{T}^4 \rightarrow \mathbb{T}^4$ is ergodic and has linear uniform growth of the derivative. Moreover, it is not metrically isomorphic to any 3-steps skew product.*

Proof. The first hypothesis of the theorem is obvious. Now suppose, contrary to our claim, that $f \times T_\varphi$ is metrically isomorphic to a 3-steps skew product. Then $f \times T_\varphi$ is measure theoretically distal (has generalized discrete spectrum in the terminology of [15]). However, $f \times T_\varphi$ has a weakly mixing factor, which is impossible, because measure theoretically distal and weakly mixing dynamical systems are disjoint (see [6]). $\square$

In the remainder of this section we present two examples of area-preserving weakly mixing diffeomorphisms with sublinear growth of the derivative.

For given $\alpha \in \mathbb{T}$ and $\beta : \mathbb{T} \rightarrow \mathbb{R}$ we will denote by $T_{\alpha,\beta} : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$ the skew product $T_{\alpha,\beta}(x_1, x_2) = (x_1 + \alpha, x_2 + \beta(x_1))$. Let $a \in \mathbb{T}$ be an irrational number and let $b : \mathbb{T} \rightarrow \mathbb{R}$ be a positive C^∞ -function with $\int_{\mathbb{T}} b(x) dx = 1$. Let σ^t denote the flow on $\mathbb{T} \times \mathbb{R}$ given by $\sigma^t(x_1, x_2) = (x_1, x_2 + t)$. Let us consider the quotient space $M_{a,b} = \mathbb{T} \times \mathbb{R} / \sim$, where the relation $\sim$ is defined by $(x_1, x_2) \sim (y_1, y_2)$ iff $(x_1, x_2) = T_{a,-b}^k(y_1, y_2)$ for an integer k . Then the quotient flow $\sigma_{a,b}^t$ of the action σ^t by the relation $\sim$ is the special flow constructed over the rotation by a and under the function b . By Lemma 2 in [3] and Theorem 1 in [10], there exists a C^∞ -diffeomorphism $\rho : M_{a,b} \rightarrow \mathbb{T}^2$ such that the flow $\varphi^t = \rho \circ \sigma_{a,b}^t \circ \rho^{-1}$ is a Hamiltonian flow on $\mathbb{T}^2$ with no fixed points, i.e. there exists C^∞ -function $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $D\xi$ is $\mathbb{Z}^2$ -periodic, non-zero at each point and

$$\frac{d}{dt}\varphi^t(\bar{x}) = \begin{bmatrix} \xi_{x_2}(\varphi^t(\bar{x})) \\ -\xi_{x_1}(\varphi^t(\bar{x})) \end{bmatrix}.$$

We will identify ρ with a diffeomorphism $\rho : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

$$\begin{aligned} \rho(x_1 + 1, x_2) &= \rho(x_1, x_2) + (N_{11}, N_{12}), \\ \rho(x_1 + a, x_2 - b(x_1)) &= \rho(x_1, x_2) + (N_{21}, N_{22}) \end{aligned}$$

for any $(x_1, x_2) \in \mathbb{R}^2$, where $N \in GL_2(\mathbb{Z})$. Then

$$(28) \quad D\rho(x_1 + 1, x_2) = D\rho(x_1, x_2)$$

$$(29) \quad D\rho(T_{a,-b}^n(x_1, x_2)) \begin{bmatrix} 1 & 0 \\ -Db^{(n)}(x_1) & 1 \end{bmatrix} = D\rho(x_1, x_2)$$

for any integer n .

Let $T_{\alpha,\beta} : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$ be a commuting with $T_{a,-b}$ skew product, where $\beta : \mathbb{T} \rightarrow \mathbb{R}$ is of class C^∞ . Then $T_{\alpha,\beta}$ can be treated as the C^∞ -diffeomorphism of $M_{a,b}$. Let us denote by $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ the area-preserving C^∞ -diffeomorphism given by $f := \rho \circ T_{\alpha,\beta} \circ \rho^{-1}$.

Lemma 12 *The diffeomorphism $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ has sublinear growth of the derivative.*

Proof. Since

$$Df^n(\bar{x}) = D\rho(T_{\alpha,\beta}^n \circ \rho^{-1}(\bar{x})) \begin{bmatrix} 1 & 0 \\ D\beta^{(n)}(\rho_1^{-1}(\bar{x})) & 1 \end{bmatrix} D\rho^{-1}(\bar{x}),$$

it suffices to show that

$$\frac{1}{n} D\rho(T_{\alpha,\beta}^n(x_1, x_2)) \begin{bmatrix} 1 & 0 \\ D\beta^{(n)}(x_1) & 1 \end{bmatrix} \rightarrow 0$$

uniformly on the set $M' = \{(x_1, x_2) : x_1 \in \mathbb{R}, 0 \leq x_2 \leq b(x_1)\}$. For every $(x_1, x_2) \in \mathbb{R}^2$ let us denote by $n(x_1, x_2)$ the unique integer number such that $T_{a,-b}^{n(x_1, x_2)}(x_1, x_2) \in M'$, i.e.

$$b^{(n(x_1, x_2))}(x_1) \leq x_2 \leq b^{(n(x_1, x_2)+1)}(x_1).$$

Let c, C be positive constants such that $0 < c \leq b(x) \leq C$ for every $x \in \mathbb{T}$. Then

$$c|n(x_1, x_2)| \leq |x_2| \leq C|n(x_1, x_2)| + C.$$

Since

$$\begin{aligned} & \frac{1}{n} D\rho(T_{\alpha, \beta}^n(x_1, x_2)) \begin{bmatrix} 1 & 0 \\ D\beta^{(n)}(x_1) & 1 \end{bmatrix} \\ &= D\rho(T_{a, -b}^{n(T_{\alpha, \beta}^n(x_1, x_2))}) T_{\alpha, \beta}^n(x_1, x_2) \frac{1}{n} \begin{bmatrix} 1 & 0 \\ -Db^{(n(T_{\alpha, \beta}^n(x_1, x_2)))}(x_1 + n\alpha) + D\beta^{(n)}(x_1) & 1 \end{bmatrix} \end{aligned}$$

(by (29)), $D\rho$ is bounded on M' (by (28)) and $n^{-1}D\beta^{(n)}$ tends uniformly to zero, it suffices to show that

$$\frac{1}{n} Db^{(n(T_{\alpha, \beta}^n(x_1, x_2)))}(x_1 + n\alpha) \rightarrow 0$$

uniformly on M' . To prove it, first observe that

$$|n(T_{\alpha, \beta}^n(x_1, x_2))| \leq c^{-1}|x_2 + \beta^{(n)}(x_1)| \leq k_1 + k_2 n,$$

for any natural n and every $(x_1, x_2) \in M'$, where $k_1 = C/c$ and $k_2 = \|\beta\|_{\sup}/c$. Fix $\varepsilon > 0$. Let n_0 be a natural number such that $|n| \geq n_0$ implies

$$\frac{1}{|n|} \|Db^{(n)}\|_{\sup} < \varepsilon/2k_2 \quad \text{and} \quad k_1 + k_2 n \leq 2k_2$$

for any integer n . Assume that n is a natural number such that $n \geq \|b\|_{C^1} n_0 / \varepsilon$. Let $(x_1, x_2) \in M'$. If $|n(T_{\alpha, \beta}^n(x_1, x_2))| \|b\|_{C^1} / n < \varepsilon$, then

$$\left| \frac{1}{n} Db^{(n(T_{\alpha, \beta}^n(x_1, x_2)))}(x_1 + n\alpha) \right| \leq \frac{|n(T_{\alpha, \beta}^n(x_1, x_2))|}{n} \|b\|_{C^1} < \varepsilon.$$

Otherwise,

$$|n(T_{\alpha, \beta}^n(x_1, x_2))| \geq \varepsilon n / \|b\|_{C^1} \geq n_0.$$

Then

$$\begin{aligned} & \left| \frac{1}{n} Db^{(n(T_{\alpha, \beta}^n(x_1, x_2)))}(x_1 + n\alpha) \right| \\ & \leq \frac{|n(T_{\alpha, \beta}^n(x_1, x_2))|}{n} \frac{1}{|n(T_{\alpha, \beta}^n(x_1, x_2))|} \|Db^{(n(T_{\alpha, \beta}^n(x_1, x_2)))}\|_{\sup} \\ & < \frac{k_1 + k_2 n}{n} \varepsilon / 2k_2 \leq \varepsilon, \end{aligned}$$

which completes the proof. $\square$

Proposition 2 (see [1]) *For every C^2 -function $\beta : \mathbb{T} \rightarrow \mathbb{R}$ with zero mean, which is not a trigonometric polynomial there exists a dense G_δ set of irrational numbers $\alpha \in \mathbb{T}$ such that the corresponding skew product $T_{\alpha, \beta} : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$ is ergodic.*

From the proof of the Main Theorem in [14] and the nature of the weak mixing property, one can obtain the following

Proposition 3 *For every positive real analytic function $b : \mathbb{T} \rightarrow \mathbb{R}$ with $\int_{\mathbb{T}} b(x)dx = 1$, which is not a trigonometric polynomial there exists a dense G_δ set of irrational numbers $a \in \mathbb{T}$ such that the corresponding special flow $\sigma_{a,b}^t$ is weakly mixing.*

Example 1. Suppose that $\sigma_{a,b}^t$ is a weakly mixing special flow whose roof function is real analytic. Let φ^t be a Hamiltonian flow on $\mathbb{T}^2$ that is C^∞ -conjugate to the special flow $\sigma_{a,b}^t$. Then the area-preserving diffeomorphism $\varphi^1 : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ is weakly mixing and has sublinear growth of the derivative, by Lemma 12.

Example 2. By Propositions 2 and 3, there exist a C^∞ -function $\beta : \mathbb{T} \rightarrow \mathbb{R}$ with zero mean and an irrational numbers $\alpha \in \mathbb{T}$ such that the corresponding skew product $T_{\alpha,\beta} : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$ is ergodic and for no real $r \neq 0$ there exist $c \in \mathbb{T}$ and a measurable function $c_r : \mathbb{T} \rightarrow \mathbb{T}$ satisfying

$$c_r(x + \alpha)e^{2\pi i r \beta(x)} = c \cdot c_r(x).$$

Assume that $b : \mathbb{T} \rightarrow \mathbb{R}$ is a positive C^∞ -function with $\int_{\mathbb{T}} b(x)dx = 1$ and $a \in \mathbb{T}$ is an irrational numbers such that the skew products $T_{\alpha,\beta}$ and $T_{a,-b}$ commute and $T_{a,-b+1}$ is not measurably isomorphic to any iteration of $T_{\alpha,\beta}$. Let us consider the special flow $\sigma_{a,b}^t$ on $M_{a,b}$ and the diffeomorphism $T_{\alpha,\beta} : M_{a,b} \rightarrow M_{a,b}$. Then $T_{\alpha,\beta} : M_{a,b} \rightarrow M_{a,b}$ is ergodic, by the ergodicity of $T_{\alpha,\beta} : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$. Moreover, $T_{\alpha,\beta} : M_{a,b} \rightarrow M_{a,b}$ is weakly mixing. Indeed, suppose, contrary to our claim, that $T_{\alpha,\beta}$ has an eigenvalue $c \in \mathbb{T}$. Then there exists a measurable function $F : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{T}$ such that

$$(30) \quad F \circ T_{\alpha,\beta} = cF \quad \text{and} \quad F \circ T_{a,-b} = F.$$

It follows that

$$F \circ \sigma^t \circ T_{\alpha,\beta} = F \circ T_{\alpha,\beta} \circ \sigma^t = cF \circ \sigma^t,$$

and hence that

$$F\sigma^t / F \circ T_{\alpha,\beta} = F\sigma^t / F$$

for any real t . By the ergodicity of $T_{\alpha,\beta}$, there exists a measurable function $c : \mathbb{R} \rightarrow \mathbb{T}$ such that $F\sigma^t = c(t)F$. Then

$$c(t+s)F = F\sigma^{t+s} = F\sigma^t\sigma^s = c(t)c(s)F$$

for all real t, s . Therefore there exists a real number r such that $c(t) = e^{2\pi i r t}$. Consequently,

$$F(x_1, x_2) = F(x_1)e^{2\pi i r x_2},$$

where $F : \mathbb{T} \rightarrow \mathbb{T}$ is given by $F(x) = F(x, 0)$. From (30), we conclude that

$$F(x_1 + \alpha)e^{2\pi i r (x_2 + \beta(x_1))} = cF(x_1)e^{2\pi i r x_2},$$

and finally that

$$F(x_1 + \alpha)e^{2\pi i r \beta(x_1)} = cF(x_1),$$

which contradicts our assumption.

Let φ^t be a Hamiltonian flow on $\mathbb{T}^2$ that is C^∞ -conjugate to the special flow $\sigma_{a,b}^t$, via a C^∞ -diffeomorphism $\rho : M_{a,b} \rightarrow \mathbb{T}^2$. Let us consider the area-preserving diffeomorphism $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ given by $f := \rho \circ T_{\alpha,\beta} \circ \rho^{-1}$. Then f is weakly mixing and has sublinear growth of the derivative, by Lemma 12.

A

For given $\alpha \in \mathbb{R}$, $\varphi : \mathbb{T} \rightarrow \mathbb{T}$ and $\varepsilon = \pm 1$ let $T_{\alpha,\varphi,\varepsilon} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ denote the diffeomorphism $T_{\alpha,\varphi,\varepsilon}(x_1, x_2) = (x_1 + \alpha, \varepsilon x_2 + \varphi(x_1))$.

Theorem 13 *Let $f : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be an area-preserving C^2 -diffeomorphism. Suppose that there exist an irrational number α and a C^2 -function $\xi : \mathbb{T}^2 \rightarrow \mathbb{T}$ such that*

- $\xi \circ f = \xi + \alpha$,
- $0 < c \leq \|D\xi(\bar{x})\|$ for any $\bar{x} \in \mathbb{T}^2$,
- $(d_1(\xi), d_2(\xi)) = (p_1, p_2) \neq (0, 0)$, where p_1, p_2 are relatively prime integer numbers.

Then there exist an area-preserving C^2 -diffeomorphism $\psi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ and a C^2 -cocycle $\varphi : \mathbb{T} \rightarrow \mathbb{T}$ such that

$$f \circ \psi = \psi \circ T_{\alpha,\varphi,\varepsilon}$$

where $\varepsilon = \det Df$ and $\xi \circ \psi(x_1, x_2) = x_1$.

Proof. Without restriction of generality we can assume that $\xi(0, 0) = 0$. For every $s \in \mathbb{R}$ set

$$A_s = \{\bar{x} \in \mathbb{R}^2 : \xi(\bar{x}) = s\}.$$

Since $0 < c \leq \|D\xi(\bar{x})\|$ for any $\bar{x} \in \mathbb{R}^2$, we see that A_s is a curve. Moreover,

$$(31) \quad (x_1, x_2) \in A_s \Rightarrow (x_1 + p_2, x_2 - p_1) \in A_s \quad \text{and} \quad \bar{x} \in A_s \Rightarrow f(\bar{x}) \in A_{s+\alpha}.$$

Let q_1, q_2 be integer numbers such that $p_1 q_1 + p_2 q_2 = 1$. Then

$$(x_1, x_2) \in A_s \Rightarrow (x_1 + q_1, x_2 + q_2) \in A_{s+1}.$$

Let $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$ be a C^2 -function such that:

- $\gamma(0) = (0, 0)$,
- $\gamma(s+1) = \gamma(s) + (q_1, q_2)$ for any real s ,




Figure 1.

- $\xi \circ \gamma(s) = s$ for any real s (see Fig. 1).

For every $s \in \mathbb{R}$, denote by $\psi(s, \cdot) : \mathbb{R} \rightarrow \mathbb{R}^2$ the solution of the following differential equation

$$\begin{aligned} \frac{d}{dt}\psi(s, t) &= \begin{bmatrix} -\xi_{x_2}(\psi(s, t)) \\ \xi_{x_1}(\psi(s, t)) \end{bmatrix} \\ \psi(s, 0) &= \gamma(s). \end{aligned}$$

Clearly, $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is of class C^2 and

$$\psi(s+1, t) = \psi(s, t) + (q_1, q_2)$$

for any $(s, t) \in \mathbb{R}^2$. Moreover,

$$\frac{d}{dt}\xi \circ \psi(s, t) = D\xi(\psi(s, t)) \cdot \frac{d}{dt}\psi(s, t) = 0.$$

Hence

$$(32) \quad \xi(\psi(s, t)) = \xi(\psi(s, 0)) = \xi(\gamma(s)) = s.$$

It follows that

$$(33) \quad A_s = \{\psi(s, t) : t \in \mathbb{R}\},$$

because $\|\frac{d}{dt}\psi(s, t)\| \geq c$. Therefore ψ is a surjection. Moreover,

$$\begin{aligned} (34) \quad \det D\psi(s, t) &= \det \begin{bmatrix} \frac{d}{ds}\psi_1(s, t) & -\xi_{x_2}(\psi(s, t)) \\ \frac{d}{ds}\psi_2(s, t) & \xi_{x_1}(\psi(s, t)) \end{bmatrix} \\ &= D\xi(\psi(s, t)) \frac{d}{ds}\psi(s, t) = \frac{d}{ds}\xi \circ \psi(s, t) = 1. \end{aligned}$$

Next note that ψ is an injection. Suppose that $(s_1, t_1) \neq (s_2, t_2)$ and $\psi(s_1, t_1) = \psi(s_2, t_2)$. Then

$$s_1 = \xi \circ \psi(s_1, t_1) = \xi \circ \psi(s_2, t_2) = s_2,$$

by (32). Hence $t_1 \neq t_2$ and $\psi(s_1, t_1) = \psi(s_1, t_2)$. It follows that $\psi(s_1, t + t_1 - t_2) = \psi(s_1, t)$ for any real t , by the definition of ψ . Therefore, the curve A_{s_1} is close, contrary to (31). Hence ψ is an area-preserving C^2 -diffeomorphism of $\mathbb{R}^2$.

Since

$$\xi(f \circ \psi(s, t)) = \xi(\psi(s, t)) + \alpha = s + \alpha,$$

(33) shows that $f \circ \psi(s, t) = \psi(s + \alpha, \eta(s, t))$, where $\eta : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a C^2 -function. By assumption, $D\xi \circ f \cdot Df = D\xi$. It follows that

$$\begin{bmatrix} -\xi_{x_2} \circ f \\ \xi_{x_1} \circ f \end{bmatrix} = \varepsilon Df \begin{bmatrix} -\xi_{x_2} \\ \xi_{x_1} \end{bmatrix},$$

Therefore

$$\begin{aligned} \frac{d}{dt} f \circ \psi(s, t) &= Df(\psi(s, t)) \frac{d}{dt} \psi(s, t) = Df(\psi(s, t)) \begin{bmatrix} -\xi_{x_2}(\psi(s, t)) \\ \xi_{x_1}(\psi(s, t)) \end{bmatrix} \\ &= \varepsilon \begin{bmatrix} -\xi_{x_2}(f \circ \psi(s, t)) \\ \xi_{x_1}(f \circ \psi(s, t)) \end{bmatrix} = \varepsilon \begin{bmatrix} -\xi_{x_2}(\psi(s + \alpha, \eta(s, t))) \\ \xi_{x_1}(\psi(s + \alpha, \eta(s, t))) \end{bmatrix} \\ &= \varepsilon \psi_{x_2}(s + \alpha, \eta(s, t)). \end{aligned}$$

On the other hand

$$\frac{d}{dt} f \circ \psi(s, t) = \frac{d}{dt} \psi(s + \alpha, \eta(s, t)) = \psi_{x_2}(s + \alpha, \eta(s, t)) \frac{d}{dt} \eta(s, t).$$

Hence $\eta(s, t) = \varepsilon t + \eta(s, 0)$. Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be given by $\varphi(s) = \eta(s, 0)$. Then

$$f \circ \psi(s, t) = \psi(s + \alpha, \varepsilon t + \varphi(s)).$$

We only need to show that ψ is a diffeomorphism of the torus $\mathbb{T}^2$. From (31) and (33), for every real s there exists a unique real number $\tau(s) \neq 0$ such that

$$\psi(s, \tau(s)) = \psi(s, 0) + (p_2, -p_1).$$

Then the function $\tau : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, $\tau(s + 1) = \tau(s)$ and

$$\psi(s, t + \tau(s)) = \psi(s, t) + (p_2, -p_1)$$

for all real s, t , by the definition of ψ . Let $M \in GL_2(\mathbb{Z})$ denote the linear part of f . As $\xi \circ f = \xi + \alpha$, we have

$$\begin{bmatrix} p_1 & p_2 \end{bmatrix} M = \begin{bmatrix} p_1 & p_2 \end{bmatrix}.$$

It follows that

$$M \begin{bmatrix} p_2 \\ -p_1 \end{bmatrix} = \bar{\varepsilon} \begin{bmatrix} p_2 \\ -p_1 \end{bmatrix},$$

where $\bar{\varepsilon} = \det M = \pm 1$. Hence

$$f((x_1, x_2) + (p_2, -p_1)) = f(x_1, x_2) + \bar{\varepsilon}(p_2, -p_1)$$

for any $(x_1, x_2) \in \mathbb{R}^2$. Therefore

$$\begin{aligned} \psi(s + \alpha, \tau(s) + \varphi(s)) &= f \circ \psi(s, \varepsilon\tau(s)) = f(\psi(s, 0) + \varepsilon(p_2, -p_1)) \\ &= f \circ \psi(s, 0) + \varepsilon\bar{\varepsilon}(p_2, -p_1) = \psi(s + \alpha, \varphi(s)) + \varepsilon\bar{\varepsilon}(p_2, -p_1) \\ &= \psi(s + \alpha, \varepsilon\bar{\varepsilon}\tau(s + \alpha) + \varphi(s)). \end{aligned}$$

Therefore $\tau(s + 2\alpha) = \tau(s)$ for any real s . Since α is irrational, τ is continuous and periodic of period 1, we see that τ is constant. Hence

$$\psi(s + 1, t) = \psi(s, t) + (q_1, q_2),$$

$$\psi(s, t + \tau) = \psi(s, t) + (p_2, -p_1)$$

for any $(s, t) \in \mathbb{R}^2$. Let $\psi_\tau : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be given by $\psi_\tau(s, t) = \psi(s, \tau t)$. Then ψ_τ is a diffeomorphism of $\mathbb{T}^2$. We conclude from (34) that $\det D\psi_\tau(s, t) = \tau$ for all $(s, t) \in \mathbb{R}^2$, hence that $\tau = \pm 1$. It follows that

$$\psi(s + 1, t) = \psi(s, t) + (q_1, q_2),$$

$$\psi(s, t + 1) = \psi(s, t) \pm (p_2, -p_1)$$

and finally that ψ is a C^2 -diffeomorphism of $\mathbb{T}^2$ such that

$$f \circ \psi = \psi \circ T_{\alpha, \varphi, \varepsilon}. \quad \square$$

B

Let $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a continuous function and let $\bar{a}, \bar{b} \in \mathbb{R}^3$ be linearly independent vectors. Let $\bar{c} \in \mathbb{R}^3$ be a non-zero vector orthogonal to both $\bar{a}$ and $\bar{b}$.

Lemma 14 *Suppose that there exists a vector $\bar{d} \in \mathbb{R}^3$ such that*

$$h(x_1 + \bar{a}\bar{m}^T, x_2 + \bar{b}\bar{m}^T) = h(x_1, x_2) + \bar{d}\bar{m}^T$$

for all $\bar{m} \in \mathbb{Z}^3$. Then there exist $k_1, k_2 \in \mathbb{R}$ such that $\bar{d} = k_1\bar{a} + k_2\bar{b}$ and the function $\tilde{h}(x_1, x_2) = h(x_1, x_2) - k_1x_1 - k_2x_2$ is $(\bar{a}\bar{m}^T, \bar{b}\bar{m}^T)$ -periodic for all $\bar{m} \in \mathbb{Z}^3$. Moreover,

- *if $\text{rank } G(\bar{c})=0$, then $\tilde{h}$ is constant;*
- *if $\text{rank } G(\bar{c})=1$, then there exist $l_1, l_2 \in \mathbb{R}$ and a continuous function $\rho : \mathbb{T} \rightarrow \mathbb{R}$ such that $\tilde{h}(x_1, x_2) = \rho(l_1x_1 + l_2x_2)$ and $l_1\bar{a} + l_2\bar{b} \in \mathbb{Z}^3$ generates $G(\bar{c})$;*

- if $\text{rank } G(\bar{c})=2$, then $\bar{c} \in c\mathbb{Z}^3$ where $c \neq 0$.

Proof. Let us consider the function $H : \mathbb{R}^3 \rightarrow \mathbb{R}$ given by $H(\bar{x}) := h(\bar{a}\bar{x}^T, \bar{b}\bar{x}^T)$ for all $\bar{x} \in \mathbb{R}^3$. Then $H(\bar{x} + \bar{m}) = H(\bar{x}) + \bar{d}\bar{m}^T$ for all $\bar{m} \in \mathbb{Z}^3$ and

$$H((x_1, x_2, x_3)(A^{-1})^T) = h(\bar{a}A^{-1}\bar{x}^T, \bar{b}A^{-1}\bar{x}^T) = h(x_1, x_2),$$

where

$$A = \begin{bmatrix} \bar{a} \\ \bar{b} \\ \bar{c} \end{bmatrix}.$$

It follows that the function $\tilde{H} : \mathbb{R}^3 \rightarrow \mathbb{R}$ given by $\tilde{H}(\bar{x}) = H(\bar{x}) - \bar{d}\bar{x}^T$ is $\mathbb{Z}^3$ -periodic. Let $A^{-1} = [\bar{a}^T \ \bar{b}^T \ \bar{c}^T]$. Then

$$\bar{d}\bar{a}'^T x_1 + \bar{d}\bar{b}'^T x_2 + \bar{d}\bar{c}'^T x_3 = h(x_1, x_2) - \tilde{H}((x_1, x_2, x_3)(A^{-1})^T).$$

As $\tilde{H}$ is bounded, we have $\bar{d}\bar{c}'^T = 0$. Define

$$\tilde{h}(x_1, x_2) := h(x_1, x_2) - k_1 x_1 - k_2 x_2,$$

where $k_1 = \bar{d}\bar{a}'^T$ and $k_2 = \bar{d}\bar{b}'^T$. Then $k_1\bar{a} + k_2\bar{b} = \bar{d}$ and

$$\tilde{h}(x_1, x_2) = \tilde{H}((x_1, x_2, x_3)(A^{-1})^T)$$

for all $(x_1, x_2, x_3) \in \mathbb{R}^3$. Therefore

$$\begin{aligned} \tilde{h}(x_1 + \bar{a}\bar{m}^T, x_2 + \bar{b}\bar{m}^T) &= \tilde{H}(((x_1, x_2, x_3) + \bar{m}A^T)(A^{-1})^T) \\ &= \tilde{H}(((x_1, x_2, x_3)(A^{-1})^T + \bar{m}) = \tilde{h}(x_1, x_2) \end{aligned}$$

for all $\bar{m} \in \mathbb{Z}^3$. Moreover,

$$\tilde{H}(x_1, x_2, x_3) = \tilde{h}(\bar{a}\bar{x}^T, \bar{b}\bar{x}^T).$$

The function $\tilde{H}$ can be represented as the Fourier series

$$\tilde{H}(\bar{x}) = \sum_{\bar{m} \in \mathbb{Z}^3} a_{\bar{m}} \exp 2\pi i \bar{m} \bar{x}^T,$$

where the series converges in $L^2(\mathbb{T}^3)$. It follows that

$$\tilde{h}(x_1, x_2) = \sum_{\bar{m} \in \mathbb{Z}^3} a_{\bar{m}} \exp 2\pi i (\bar{m}\bar{a}'^T x_1 + \bar{m}\bar{b}'^T x_2 + \bar{m}\bar{c}'^T x_3)$$

in $L^2(\mathbb{T}_A^3)$. Therefore, $\bar{m}\bar{c}'^T = 0$, wherever $a_{\bar{m}} \neq 0$. Hence

$$\tilde{H}(\bar{x}) = \sum_{\bar{m} \in G(\bar{c})} a_{\bar{m}} \exp 2\pi i \bar{m} \bar{x}^T,$$

because $G(\bar{c}) = G(\bar{c}')$.

Suppose that $\text{rank } G(\bar{c}) = 0$. Then $\tilde{H}$, and consequently $\tilde{h}$ is constant.

Now suppose that $\text{rank } G(\bar{c}) = 1$. Then $\tilde{H}(\bar{x}) = \rho(\bar{m}\bar{x}^T)$, where $\bar{m}$ is a generator of $G(\bar{c})$ and $\rho : \mathbb{T} \rightarrow \mathbb{R}$ is a continuous function. Moreover,

$$\tilde{h}(x_1, x_2) = \rho(\bar{m}A^{-1}\bar{x}^T) = \rho(\bar{m}\bar{a}'^T x_1 + \bar{m}\bar{b}'^T x_2)$$

and

$$\bar{m}\bar{a}'^T \bar{a} + \bar{m}\bar{b}'^T \bar{b} = \bar{m}A^{-1}A = \bar{m}.$$

Now assume that $\text{rank } G(\bar{c}) = 2$. Suppose that $\bar{m}, \bar{n} \in \mathbb{Z}^3$ generate $G(\bar{c})$. Then $\bar{c} = c\bar{m} \times \bar{n} \in c\mathbb{Z}^3$. $\square$

Lemma 15 *Let $\bar{c} \in \mathbb{Z}^3 \setminus \{0\}$. Then there exists a pair of generators $\bar{a}, \bar{b} \in \mathbb{Z}^3$ of $G(\bar{c})$ such that $\Lambda(\bar{a}, \bar{b}) = \{(\bar{a}\bar{m}^T, \bar{b}\bar{m}^T) \in \mathbb{Z}^2 : \bar{m} \in \mathbb{Z}^3\} = \mathbb{Z}^2$.*

Proof. Our proof starts with the observation that we can assume that c_1, c_2, c_3 are relatively prime. Let $p_1 := \gcd(c_2, c_3)$, $p_2 := \gcd(c_3, c_1)$, $p_3 := \gcd(c_1, c_2)$. Then p_1, p_2, p_3 are pairwise relatively prime and there exist pairwise relatively prime integer numbers k_1, k_2, k_3 such that $c_1 = k_1 p_2 p_3$, $c_2 = p_1 k_2 p_3$, $c_3 = p_1 p_2 k_3$. Let us choose integer numbers x, y such that $k_2 x + k_3 y = 1$. Then

$$\bar{a} := (p_1, -p_2 k_1 x, -p_3 k_1 y) \quad \text{and} \quad \bar{b} := (0, -p_2 k_3, p_3 k_2)$$

generates the group $G(\bar{c})$. Indeed, suppose that $(m_1, m_2, m_3) \in G(\bar{c})$. Since $m_1 c_1 = -m_2 c_2 - m_3 c_3$ we see that $p_1 | m_1 c_1$, hence that $p_1 | m_1$, because c_1 and p_1 are relatively prime. Similarly, there exist $n_1, n_2, n_3 \in \mathbb{Z}$ such that $m_1 = p_1 n_1$, $m_2 = p_2 n_2$, $m_3 = p_3 n_3$. As $(m_1, m_2, m_3) \in G(\bar{c})$ we have

$$n_1 k_1 + n_2 k_2 + n_3 k_3 = 0.$$

Since $n_1 k_1 + n_1(-k_1 x)k_2 + n_1(-k_1 y)k_3 = 0$ we obtain

$$(n_2 + n_1 k_1 x)k_2 + (n_3 + n_1 k_1 y)k_3 = 0.$$

Therefore there exists $s \in \mathbb{Z}$ such that $n_2 + n_1 k_1 x = -s k_3$ and $n_3 + n_1 k_1 y = s k_2$. It follows that

$$(n_1, n_2, n_3) = n_1(1, -k_1 x, -k_1 y) + s(0, -k_3, k_2),$$

hence that $(m_1, m_2, m_3) = n_1 \bar{a} + s \bar{b}$. Let r, t be integer numbers such that $r p_1 - t c_1 = 1$. Then

$$(\bar{a}, \bar{b})(r, t p_3 k_2, t p_2 k_3)^T = (1, 0).$$

Moreover

$$(\bar{a}, \bar{b})(0, r', t')^T = (s', 1),$$

whenever $-p_2 k_3 r' + p_3 k_2 t' = 1$. It follows that $(1, 0)$ and $(s', 1)$ belong to $\Lambda(\bar{a}, \bar{b})$, and finally that $\Lambda(\bar{a}, \bar{b}) = \mathbb{Z}^2$. $\square$

Corollary 1 *For any pair of generators $\bar{a}', \bar{b}' \in \mathbb{Z}^3$ of $G(\bar{c})$ we have $\Lambda(\bar{a}', \bar{b}') = \mathbb{Z}^2$.*

Proof. Suppose that $\bar{a}', \bar{b}' \in \mathbb{Z}^3$ generate $G(\bar{c})$. Then there exists $K \in GL_2(\mathbb{Z})$ such that $(\bar{a}'^T, \bar{b}'^T) = (\bar{a}^T, \bar{b}^T)K$. Therefore

$$\Lambda(\bar{a}', \bar{b}') = \Lambda(\bar{a}, \bar{b})K = \mathbb{Z}^2. \quad \square$$

References

- [1] J. Aaronson, M. Lemańczyk, C. Mauduit, H. Nakada, *Koksma's inequality and group extensions of Kroneker transformations*, in: Algorithms, Fractals and Dynamics (Okayama and Kyoto, 1992), Y. Takahashi (ed.), Plenum, New York, 1995, 27-50.
- [2] I.P. Cornfeld, S.W. Fomin, J.G. Sinai, *Ergodic Theory*, Springer, Berlin, 1982.
- [3] B. Fayad, A. Katok, A. Windsor, *Mixed spectrum reparameterizations of linear flows on $\mathbb{T}^2$* , Mosc. Math. J. **1** (2001), 521-537.
- [4] K. Frączek, *Linear growth of the derivative for measure-preserving diffeomorphism*, Coll. Math. **84/85** Part 1 (2000), 147-157.
- [5] K. Frączek, *Measure-preserving diffeomorphisms of the torus*, Ergod. Th. & Dynam. Sys. **21** (2001), 1759-1781.
- [6] H. Furstenberg, *Disjointness in ergodic theory, minimal sets, and a problem in diophantine approximation*, Math. Systems Theory **1** (1967), 1-49.
- [7] M. Herman, *Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations*, Publ. Mat. IHES **49** (1979), 5-234.
- [8] A. Iwanik, M. Lemańczyk, D. Rudolph, *Absolutely continuous cocycles over irrational rotations*, Isr. J. Math. **83** (1993), 73-95.
- [9] A. Manning, *There are no new Anosov diffeomorphisms on tori*, Amer. J. Math. **96** (1974), 422-429.
- [10] J. Moser, *On the volume elements on a manifold*, Trans. Amer. Math. Soc. **120** (1965), 286-294.
- [11] L. Polterovich, J.-C. Sikorav, *A linear isoperimetric inequality for the punctured Euclidean plane*, preprint math.GR/0106216.
- [12] L. Polterovich, *Growth of maps, distortion in groups and symplectic geometry*, preprint math.DS/0111050.
- [13] K. Schmidt, *On Recurrence*, Z. Wahrsch. Verw. Gebiete **68** (1984), 75-95.

- [14] M.D. Shklover, *On classical dynamical systems on the torus with continuous spectrum*, Izv. Vyssh. Uchebn. Zaved., Mat. **10** (1967), 113-124.
- [15] R.J. Zimmer, *Ergodic actions with generalized discrete spectrum*, Illinois J. Math. **20** (1975), 555-588.

Faculty of Mathematics and Computer Science,
Nicholas Copernicus University
ul. Chopina 12/18
87-100 Toruń, Poland

Institute of Mathematics
Polish Academy of Science
ul. Chopina 12
87-100 Toruń, Poland

E-mail: fraczek@mat.uni.torun.pl